

Classical electrodynamics of point charges

Massimo Marino*

*INFN - Dipartimento di Fisica, Università di Milano,
via Celoria 16, I-20133 Milano, Italy*

Abstract

A simple mathematical procedure is introduced which allows redefining in an exact way divergent integrals and limits that appear in the basic equations of classical electrodynamics with point charges. In this way all divergences are at once removed without affecting the locality and the relativistic covariance of the theory, and with no need for mass renormalization. The procedure is first used to obtain a finite expression for the electromagnetic energy-momentum of the system. We show that the relativistic Lorentz-Dirac equation can be deduced from the conservation of this electromagnetic energy-momentum plus the usual mechanical term. Then we derive a finite lagrangian, which depends on the particle variables and on the actual electromagnetic potentials at a given time. From this lagrangian the equations of motion of both particles and fields can be derived via Hamilton's variational principle. The hamiltonian formulation of the theory can be obtained in a straightforward way. This leads to an interesting comparison between the resulting divergence-free expression of the hamiltonian functional and the standard renormalization rules for perturbative quantum electrodynamics.

03.50.De, 11.10.Ef

Typeset using REVTeX

*Email address: massimo.marino@mi.infn.it

I. INTRODUCTION

The relativistic Lorentz-Dirac equation was derived in 1938 [1] to describe the motion of a classical point charge in an arbitrarily assigned external field, taking into account its electromagnetic self-interaction and therefore the effect of radiation reaction (useful reviews on this subject can be found in Ref. [2,3]).

Dirac believed that, once a finite classical theory for point-like charges had been obtained, then by applying appropriate rules of canonical quantization it should be possible to formulate the quantum theory in a way that avoided the occurrence of troublesome infinities. Already in the final part of his original paper [1], with the explicit purpose of suggesting the path towards a possible quantization, Dirac showed that his theory could be deduced from a variational principle, by making use of a non-local action which had been previously proposed by Fokker [4] to describe a system of charged particles interacting at a distance through the average of the retarded and advanced fields. However Dirac soon realized that such an action could not allow to reexpress the dynamics in hamiltonian form. For this reason, soon thereafter he produced a second paper [5] specifically devoted to the problem of achieving a canonical quantization of his theory. In this work he abandoned the action principle, and attempted to write down directly a four-dimensional hamiltonian in terms of an unphysical field dependent on the proper time of each particle constituting the system under consideration. For this field Dirac skilfully postulated particular *ad hoc* Poisson brackets, involving an arbitrary time-like constant four-vector λ , in such a way that his covariant equation of motion for the particles was recovered in the limit of vanishing λ . It is not surprising that such an involute approach could hardly lend itself to further generalizations, and Dirac was unable to use it as the starting point for a consistent formulation of quantum electrodynamics. It is also interesting to observe that, although Dirac had derived in [1] his equation of motion using essentially the principle of local energy-momentum conservation, he did not provide a general expression for the global conserved quantities. Besides, the conservation laws no longer emerged in an obvious way from the canonical formalism that he proposed.

With the advent of renormalization theory, perturbative quantum electrodynamics developed in an impressive way without relying on any input from classical theory for the formal treatment of divergences. However, research on the classical theory was never abandoned. In particular, a new finite classical action describing the behavior of both particles and fields was obtained by Rohrlich [2,6] by adding an appropriate term to the original expression of Fokker and Dirac. However, this result could not help in providing a canonical formulation of the theory. Furthermore, in the action the electromagnetic field was split as the sum of a source-free term plus the contribution generated by the particles through the retarded and advanced potentials. This contrasted with the lagrangian formulation of QED which, despite the subtleties related to the problem of providing an adequate renormalization, is always expressed only in terms of the full interacting electromagnetic field. Similar considerations apply also for other studies subsequently appeared, in which the expression of the electromagnetic energy-momentum in terms of the retarded fields was subjected to an accurate analysis [7], and a different kind of lagrangian, based on the distribution theory, was derived [8].

In the present paper we shall propose a simple and general method to evaluate the inte-

grals of the electromagnetic energy-momentum, lagrangian and hamiltonian densities over spatial regions containing the point particles. In a similar way, we shall assign a finite value to the fields and potentials on the particles' world-lines, by generalizing a procedure previously proposed by Teitelboim and Lopez [9,10]. In this way practically all the familiar formulae of electrodynamics, which are commonly applied when the particles' self-interaction is neglected, can be immediately generalized to the self-interacting case by appropriately reinterpreting each individual term according to the given prescriptions. The resulting equations of motion will be the relativistic Lorentz-Dirac equation for the particles and the Maxwell equations with point-like sources for the fields. In particular we shall obtain:

1. A finite expression for the total energy-momentum of the system of fields and particles, which is Lorentz-covariant and conserved on the solutions of the equations of motion;
2. A finite lagrangian which determines the dynamics of both fields and particles according to Hamilton's variational principle;
3. A finite hamiltonian and a system of Poisson's brackets which express the dynamics in canonical form.

In the treatment here presented, the infinities will be consistently removed without altering any of the fundamental formal properties of the theory, including locality and relativistic covariance. A remarkable property of this procedure is that it allows expressing the four-momentum, the lagrangian and the hamiltonian of the system only in terms of the particle variables and of the total instantaneous electromagnetic field, i.e. without any unphysical splitting into retarded, advanced, incoming or outgoing parts. This will provide a formulation of the theory which is in complete agreement with the requirements of local relativistic field-theory. Furthermore, no diverging parameter will have to be introduced, since only the physical mass of the particles will be used. Due to the close correspondence with the quantum theory, we shall show that these results may also suggest a new way of formulating the renormalization procedure of quantum electrodynamics.

The paper is organized as follows. In Sec. II we define the renormalized energy-momentum associated with the electromagnetic field in the presence of point charges, and we discuss how the conservation of the sum of this energy-momentum and the familiar mechanical term associated with the particles (with mass parameter equal to the observable mass m) determines the equation of motion of the particles, provided that the field satisfies the usual Maxwell equations. Actually, this provides a new rigorous and straightforward derivation of the relativistic Lorentz-Dirac equation. We shall comment on the mathematical structure of the system of coupled equations of motion for particles and fields, and on the non-trivial constraints that are to be imposed on the initial data of the corresponding Cauchy problem. In Sec. III we define the renormalized lagrangian of the system, and we show how the equations of motion of both particles and fields can be derived from an action principle. We shall see that on the solutions of the field equations the value assumed by our lagrangian can be approximated by an expression corresponding to an electron of finite radius r and bare mass $m_0 = m - q^2/8\pi r$, although the canonical forms of the two theories remain essentially inequivalent even in the limit $r \rightarrow 0$. In Sec. IV we give a precise mathematical definition for the partial derivatives of the lagrangian with respect to the particle

variables, and we make use of this definition to put the equations of motion in the standard form of Euler-Lagrange equations. The reformulation of the theory in the hamiltonian formalism will then be obtained in a straightforward way via a Legendre transformation. In Sec. V we obtain an equivalent expression for the renormalized hamiltonian in terms of the Fourier transform of the electromagnetic potential, and in Sec. VI we make use of this expression to illustrate the close relationship with the renormalization rules for the electron self-energy in perturbative quantum electrodynamics.

Most of the mathematical proofs and technical developments are worked out in the Appendixes. In particular, in App. A we consider the retarded and advanced fields near a point charge in arbitrary motion, and we show that they can be asymptotically expanded as the sum of a series of covariant homogeneous functions of the space-time variables. This essential result will be extensively applied throughout the paper.

II. THE RENORMALIZED ENERGY-MOMENTUM

A. The electromagnetic energy-momentum in the presence of point charges

In the elementary case of electrostatics, the energy of the electric field associated with a non-singular spatially confined charge distribution $\rho(\mathbf{x})$ is given by

$$U = \frac{1}{2} \int d^3\mathbf{x} \mathbf{E}^2(\mathbf{x}), \quad (2.1)$$

where

$$\mathbf{E}(\mathbf{x}) = \frac{1}{4\pi} \int d^3\mathbf{x}' \rho(\mathbf{x}') \frac{\mathbf{x} - \mathbf{x}'}{|\mathbf{x} - \mathbf{x}'|^3}. \quad (2.2)$$

When the charge distribution is constituted by a set of N point charges $q_1, q_2, \dots, q_N$ located at the points $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_N$, Eq. (2.2) takes the form

$$\mathbf{E}(\mathbf{x}) = \frac{1}{4\pi} \sum_{n=1}^N q_n \frac{\mathbf{x} - \mathbf{z}_n}{|\mathbf{x} - \mathbf{z}_n|^3}.$$

It is well known that in such a case the integral on the right-hand side of Eq. (2.1) becomes divergent. However, introducing a cut-off at a distance r from the point charges, one can define

$$\overline{U}(r) \equiv \frac{1}{2} \int d^3\mathbf{x} \mathbf{E}^2(\mathbf{x}) \prod_{n=1}^N \theta(|\mathbf{x} - \mathbf{z}_n| - r),$$

where θ is the usual step function. Then an elementary integration provides

$$\overline{U}(r) = \frac{1}{8\pi r} \sum_{n=1}^N q_n^2 + \frac{1}{4\pi} \sum_{n < m} \frac{q_n q_m}{|\mathbf{z}_n - \mathbf{z}_m|} + O(r) \quad \text{for } r \rightarrow 0. \quad (2.3)$$

It can be seen therefore that the divergent part is only represented by an irrelevant constant term which can be safely dropped, leaving us with the familiar Coulomb potential energy:

$$U(\mathbf{z}_1, \dots, \mathbf{z}_N) = \frac{1}{4\pi} \sum_{n < m} \frac{q_n q_m}{|\mathbf{z}_n - \mathbf{z}_m|}. \quad (2.4)$$

The above considerations illustrate the obvious fact that the singularities associated with point charges are completely harmless in the case of electrostatics, since the physics can be completely described by finite quantities such as (2.4). In the present paper we shall show the remarkable fact that an analogous result actually holds also for relativistic electrodynamics. To this end, it is useful to introduce a general limiting procedure for functions near a first-order singular point.

Definition 1 *Given a function $f(r)$ of a variable r , having in a neighborhood of $r = 0$ a behavior of the form*

$$f(r) = \frac{a}{r} + b + O(r),$$

we define its “ $\mathcal{R}$ -limit” (henceforth indicated by the symbol $\mathcal{R} \lim$) for $r \rightarrow 0$ as

$$\mathcal{R} \lim_{r \rightarrow 0} f(r) \equiv \lim_{r \rightarrow 0} \frac{d}{dr} [r f(r)] = b.$$

Correspondingly, given a function $G(\mathbf{x})$ of the spatial coordinates $\mathbf{x}$, having in a set V the singular points $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_N$, and a behavior such that

$$\int_V d^3 \mathbf{x} G(\mathbf{x}) \prod_{n=1}^N \theta(|\mathbf{x} - \mathbf{z}_n| - r) = \frac{a'}{r} + b' + O(r), \quad (2.5)$$

we define its “ $\mathcal{R}$ -integral” over V as

$$\mathcal{R} \int_V d^3 \mathbf{x} G(\mathbf{x}) \equiv \mathcal{R} \lim_{r \rightarrow 0} \int_V d^3 \mathbf{x} G(\mathbf{x}) \prod_{n=1}^N \theta(|\mathbf{x} - \mathbf{z}_n| - r) = b'.$$

It is clear that the $\mathcal{R}$ -integral defined above differs from the ordinary integral only in a neighborhood of the singular points. More precisely, if $V = V_1 \cup V_2$, where $V_1 \cap V_2 = \emptyset$ and V_2 contains no singularities, then

$$\mathcal{R} \int_V d^3 \mathbf{x} G(\mathbf{x}) = \mathcal{R} \int_{V_1} d^3 \mathbf{x} G(\mathbf{x}) + \int_{V_2} d^3 \mathbf{x} G(\mathbf{x}).$$

Applying the Definition 1 to the electrostatic problem considered at the beginning, we can write

$$U(\mathbf{z}_1, \dots, \mathbf{z}_N) = \frac{1}{4\pi} \sum_{n < m} \frac{q_n q_m}{|\mathbf{z}_n - \mathbf{z}_m|} = \mathcal{R} \int d^3 \mathbf{x} (1/2) \mathbf{E}^2(\mathbf{x}).$$

The generalization of the above expression to the electrodynamic case will be obtained simply by replacing in the integrand the electrostatic energy density $(1/2) \mathbf{E}^2$ with the full electromagnetic energy density $T^{00} = (1/2) (\mathbf{E}^2 + \mathbf{B}^2)$. The same procedure can then be applied also with the momentum density T^{0k} . In this way, for a given inertial reference

frame we are led to consider as the electromagnetic energy-momentum contained at the time t in a volume V the quantity

$$P_{V,\text{em}}^\mu(t) \equiv \mathcal{R} \int_V d^3\mathbf{x} T^{0\mu}(\mathbf{x}, t), \quad (2.6)$$

where

$$T^{\mu\nu} = F^{\mu\lambda} F^\nu{}_\lambda - \frac{1}{4} g^{\mu\nu} F^{\eta\lambda} F_{\eta\lambda} \quad (2.7)$$

is the familiar electromagnetic energy-momentum tensor (we put $c = 1$ and use the metric $g^{11} = g^{22} = g^{33} = +1$, $g^{00} = -1$). Of course, in order for the definition (2.6) to be meaningful, the integral on the right-hand side must have a behavior of the form (2.5). It is possible to prove that this requirement is actually satisfied as a consequence of the Maxwell equations for the electromagnetic field in the presence of point charges. In fact, the field near a charged particle can be decomposed as $F^{\mu\nu} = F_{\text{ext}}^{\mu\nu} + F_{\text{ret}}^{\mu\nu}$, where $F_{\text{ret}}^{\mu\nu}$ is the retarded field generated by the particle and $F_{\text{ext}}^{\mu\nu}$ satisfies the free-field equations in a neighborhood of the particle. Under the reasonable assumption that $F_{\text{ext}}^{\mu\nu}$ be non-singular in this neighborhood, the divergent part of the field can be expressed using the asymptotic expansion for $F_{\text{adv}}^{\mu\nu}$ and $F_{\text{ret}}^{\mu\nu}$ which is provided in App. A. This can be written as

$$F_{\text{adv/ret}}^{\mu\nu}(z + x) = (F^{\mu\nu})_{-2}(x) + (F^{\mu\nu})_{-1}(x) + (F_{\text{adv/ret}}^{\mu\nu})_0(x) + O(x), \quad (2.8)$$

with

$$(F)_{-2}(x) = \frac{q}{4\pi y^3} u \wedge x, \quad (2.9)$$

$$(F)_{-1}(x) = \frac{q}{4\pi y^3} \left[\left(1 - \frac{3x^2}{2y^2} \right) (\dot{u} \cdot x) u \wedge x - (u \cdot x) \dot{u} \wedge x - \frac{x^2}{2} u \wedge \dot{u} \right], \quad (2.10)$$

while $(F_{\text{adv/ret}})_{-2}$, given by Eqs. (A12) and (A13), is a term that is finite for $x \rightarrow 0$. In the above formulae, z is the space-time coordinate of the particle at a given proper time τ . We have also introduced the four-vectors $u = dz/d\tau$, $\dot{u} = du/d\tau$, the scalar $y = \sqrt{x^2 + (x \cdot u)^2}$, and we have used the notations $u \cdot x = u_\nu x^\nu$, $x^2 = x \cdot x = x_\nu x^\nu$, $(u \wedge x)^{\mu\nu} = u^\mu x^\nu - u^\nu x^\mu$ and similar. One can see that, since y is a homogeneous function of x of degree 1, the term $(F)_{-2}$ is homogeneous of degree -2 , and moreover it is odd under the inversion $x \rightarrow -x$ (leaving u and $\dot{u}$ unchanged). Consequently, when the function, as in Eq. (2.6), is restricted to the three-dimensional space with $x^0 = 0$, this term is odd under the space reflection $\mathbf{x} \rightarrow -\mathbf{x}$. Similarly, $(F)_{-1}$ is homogeneous of degree -1 and is even. It follows that $T^{0\mu}$ has a leading term of order x^{-4} which, when substituted to G in an integral of the form (2.5), gives rise to the term a'/r on the right-hand side. Besides, the term of $T^{0\mu}$ of order x^{-3} is odd, so it does not generate any logarithmic term in r . One can therefore conclude that the integral has indeed the required behavior.

It can be observed, using the same arguments, that the value of the $\mathcal{R}$ -integral on the right-hand side of Eq. (2.6) is unchanged if the cut-off regions near the point charges, instead of being spherical as in Eq. (2.5), are chosen with a different shape, provided that such a

shape be invariant under space inversion with respect to the particle position. Owing to the form of the expansion (2.8)–(2.10), it turns out that mathematical manipulations are simpler when surfaces of the form $y = \text{constant}$ are considered. Therefore we shall make use more often of the equivalent expression

$$P_{V,\text{em}}^\mu(t) = \mathcal{R} \lim_{r \rightarrow 0} \bar{P}_{V,\text{em}}^\mu(t, r), \quad (2.11)$$

with

$$\bar{P}_{V,\text{em}}^\mu(t, r) \equiv \int_V d^3\mathbf{x} T^{0\mu}(\mathbf{x}, t) \prod_{n=1}^N \theta(y_n - r) \quad (2.12)$$

and

$$y_n \equiv \sqrt{(x - z_n)^2 + [u_n \cdot (x - z_n)]^2}. \quad (2.13)$$

Using $z_n^0 = x^0 = t$, $\mathbf{z}_n = \mathbf{z}_n(t)$, $\mathbf{v}_n(t) = d\mathbf{z}_n/dt$, $\gamma_n = (1 - \mathbf{v}_n^2)^{-1/2}$, we can write also

$$y_n = \sqrt{(\mathbf{x} - \mathbf{z}_n)^2 + \gamma_n^2 [\mathbf{v}_n \cdot (\mathbf{x} - \mathbf{z}_n)]^2}, \quad (2.14)$$

from which it can be seen that the surfaces $y_n = r$ are just spheres relativistically contracted by a factor γ in the direction of movement. As shown in App. B, the term of order $1/r$ in $\bar{P}_{V,\text{em}}^\mu(t, r)$ can be easily calculated from the expansion (2.8). This gives

$$\bar{P}_{V,\text{em}}^k(t, r) = \frac{1}{4\pi r} \sum_{\mathbf{z}_n \in V} \frac{2}{3} q_n^2 u_n^k + P_{V,\text{em}}^k(t) + O(r), \quad (2.15)$$

$$\bar{P}_{V,\text{em}}^0(t, r) = \frac{1}{4\pi r} \sum_{\mathbf{z}_n \in V} q_n^2 \left(\frac{2}{3} u_n^0 - \frac{1}{6} \gamma_n^{-1} \right) + P_{V,\text{em}}^0(t) + O(r), \quad (2.16)$$

where the sums on the right-hand sides extend to all the particles contained in the volume V . One can observe from these expressions that the divergent terms of the electromagnetic energy-momentum do *not* transform as a four-vector under Lorentz transformations. In particular, for a single particle with $v \ll 1$, the $1/r$ terms alone would give rise to the well-known result $\bar{P}_{V,\text{em}}^k = (4/3) \bar{P}_{V,\text{em}}^0 v^k$. However, these divergent terms depend only on the particle velocities and contain no information about the interaction of the particles with each other or with incident external fields. This suggests that these terms may be dropped as it was done for the constant divergent term in Eq. (2.3). In so doing, we are led directly to the definition (2.6). In the following subsection we are going to provide a rigorous justification for this definition.

B. The conservation of the renormalized energy-momentum and the Lorentz-Dirac equation

If we denote with $P_V^\mu(t)$ the total energy-momentum contained the volume V at the time t , in a local theory we must expect that

$$\frac{d}{dt}P_V^\mu(t) = - \int_{\partial V} d\sigma n_i T^{i\mu}(\mathbf{x}, t), \quad (2.17)$$

where $\mathbf{n}$ is the outward unit vector normal to the surface element $d\sigma$. Clearly the right-hand side of the above equation represents the total flow of energy-momentum through the boundary ∂V of the volume V . We are now going to show how the condition expressed by Eq. (2.17), combined with the definition (2.6) for the electromagnetic energy-momentum, univocally determines the dynamics of the charged particles in a relativistically covariant way. However, in order to formulate the equations of motion, we first need to introduce the fields $\tilde{F}^{\mu\nu}(z_n)$, which are defined only on the world-line of the particles according to the relation

$$\tilde{F}^{\mu\nu}(z_n) \equiv \mathcal{R} \lim_{r \rightarrow 0} \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} F^{\mu\nu}(\mathbf{z}_n + \mathbf{x}, t), \quad (2.18)$$

with $t = z_n^0$. The integral in the solid angle on the right-hand side represents the average of the field on a spherical surface of radius r centered at $\mathbf{z}_n(t)$. Again, one can deduce from Eq. (2.8) and from symmetry considerations that the r dependence of the integral is indeed of the form required for the $\mathcal{R}$ -limit to be properly defined.

Theorem 1 *If one defines*

$$P_V^\mu(t) \equiv P_{V,\text{em}}^\mu(t) + \sum_{\mathbf{z}_n \in V} m_n u_n^\mu(t), \quad (2.19)$$

with $P_{V,\text{em}}^\mu(t)$ given by Eq. (2.6), then the condition expressed by Eq. (2.17) is satisfied for arbitrary V if and only if the particles obey the equation of motion

$$m_n \gamma_n \frac{du_n^\mu}{dt} = q_n u_{n,\nu} \tilde{F}^{\mu\nu}(z_n), \quad (2.20)$$

with $\tilde{F}^{\mu\nu}(z_n)$ given by Eq. (2.18).

The total energy-momentum of a system with N charged particles is obtained from Eq. (2.19) by taking as the volume V the whole space. This gives

$$P^\mu(t) = P_{\text{em}}^\mu(t) + \sum_{n=1}^N m_n u_n^\mu(t), \quad (2.21)$$

with

$$P_{\text{em}}^\mu(t) \equiv \mathcal{R} \int d^3\mathbf{x} T^{0\mu}(\mathbf{x}, t). \quad (2.22)$$

From Eq. (2.17) and from the usual assumption that the field vanishes sufficiently fast at infinity, it follows that $P^\mu(t)$ is a constant of the motion, i.e. satisfies

$$\frac{dP^\mu}{dt} = 0. \quad (2.23)$$

Furthermore, one can prove that it has the right transformation property required by special relativity.

Theorem 2 *The total energy-momentum P^μ given by Eq. (2.21) transforms as a four-vector under Lorentz transformations.*

The proofs of Theorems 1 and 2 are presented in Appendixes C and D respectively. It is important to observe that if we have a single charged particle in uniform rectilinear motion, and if the field at every point of space reduces to that produced by the particle (i.e. it is equal to the retarded or advanced fields, which are in this case coincident), then the expression $\overline{P}_{\text{em}}^\mu(t, r)$, defined by Eq. (2.12) with V extending to the whole space, is exactly equal to the $1/r$ term alone. From the definition of $\mathcal{R}$ -integral it follows that $P_{\text{em}}^\mu(t)$ vanishes, so that the total energy-momentum is given by $P^\mu = mu^\mu$. An analogous result holds for a set of uniformly moving particles at very large distances from each other. These considerations imply that the parameters m_n appearing in Eqs. (2.19)–(2.20) are to be identified with the total physical masses of the particles.

Considering the equation of motion (2.20), we see that the right-hand side has the familiar structure of the Lorentz force, but with the new finite field $\tilde{F}^{\mu\nu}$ instead of the total electromagnetic field $F^{\mu\nu}$, which of course diverges on the trajectory of the particles. Eq. (2.20) has the correct mathematical form that an equation of motion in a theory of interacting particles and fields is supposed to have. It expresses in fact the second-order time derivative of the particle's position as a function of the position, its first-order derivative and the instantaneous field configuration, since $\tilde{F}^{\mu\nu}(z_n)$ is a function of the field $F^{\mu\nu}(\mathbf{x}, t)$ at fixed t in a neighborhood of $\mathbf{z}_n(t)$. A complete description of the temporal evolution of the system is obtained by coupling with Eq. (2.20) the Maxwell equations

$$\frac{\partial \mathbf{B}(\mathbf{x}, t)}{\partial t} + \nabla \wedge \mathbf{E}(\mathbf{x}, t) = 0 \quad (2.24)$$

$$\frac{\partial \mathbf{E}(\mathbf{x}, t)}{\partial t} - \nabla \wedge \mathbf{B}(\mathbf{x}, t) = -\mathbf{j}(\mathbf{x}, t) = -\sum_{n=1}^N q_n \mathbf{v}_n(t) \delta(\mathbf{x} - \mathbf{z}_n(t)), \quad (2.25)$$

with the supplementary equations

$$\nabla \cdot \mathbf{E}(\mathbf{x}, t) = \rho(\mathbf{x}, t) = \sum_{n=1}^N q_n \delta(\mathbf{x} - \mathbf{z}_n(t)) \quad (2.26)$$

$$\nabla \cdot \mathbf{B}(\mathbf{x}, t) = 0 \quad (2.27)$$

to be regarded as usual as constraints on the initial conditions.

In order to clarify the meaning of Eq. (2.20), let us define for a given charged particle the field

$$\overline{F}^{\mu\nu} \equiv F^{\mu\nu} - \frac{1}{2} (F_{\text{adv}}^{\mu\nu} + F_{\text{ret}}^{\mu\nu}), \quad (2.28)$$

which remains finite on the world-line of the particle. Using Eqs. (2.8) and (A12) we can then write

$$F^{\mu\nu}(z+x) = (F^{\mu\nu})_{-2}(x) + (F^{\mu\nu})_{-1}(x) + (F^{\mu\nu})_0(x) + \overline{F}^{\mu\nu}(z) + O(x), \quad (2.29)$$

where $(F^{\mu\nu})_0(x)$, given by Eq. (A13), is a term that changes sign under the substitution $x \rightarrow -x$. Therefore, applying the definition (2.18) and taking into account the spatial antisymmetry of $(F)_{-2}$ and $(F)_0$, we obtain

$$\tilde{F}^{\mu\nu}(z) = \lim_{r \rightarrow 0} \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} [(F^{\mu\nu})_0(x) + \bar{F}^{\mu\nu}(z)] = \bar{F}^{\mu\nu}(z) \quad (2.30)$$

(a proof of Eq. (2.30) for the system of reference comoving with the particle was already given in Ref. [9]). Since the definition (2.28) is Lorentz covariant, Eq. (2.30) shows that $\tilde{F}^{\mu\nu}(z)$ actually transforms like a tensor. Substituting $\bar{F}^{\mu\nu}$ to $\tilde{F}^{\mu\nu}$ on the right-hand side of Eq. (2.20), one can recognize in the latter a well-known form of the relativistic Lorentz-Dirac equation. We note also that Eqs. (2.8) and (A12) are in agreement with the relation first proved by Dirac in [1]

$$\frac{1}{2} [F_{\text{ret}}^{\mu\nu}(x) - F_{\text{adv}}^{\mu\nu}(x)] \rightarrow \frac{2}{3} \frac{q}{4\pi} (u^\mu \ddot{u}^\nu - u^\nu \ddot{u}^\mu) \text{ for } x \rightarrow z ,$$

from which it follows that, after decomposing the field as $F^{\mu\nu} = F_{\text{ext}}^{\mu\nu} + F_{\text{ret}}^{\mu\nu}$, the Eq. (2.20) can be rewritten as

$$m\dot{u}^\mu = qu_\nu F_{\text{ext}}^{\mu\nu} + \frac{2}{3} \frac{q^2}{4\pi} (\ddot{u}^\mu - \dot{u}_\nu \dot{u}^\nu u^\mu), \quad (2.31)$$

where use has been made of the relations $u^2 = -1$, $0 = d(u^2)/d\tau = 2\dot{u} \cdot u$, $0 = d(\dot{u} \cdot u)/d\tau = \dot{u}^2 + \ddot{u} \cdot u$. In the familiar form (2.31) the Lorentz-Dirac equation appears explicitly as a *third-order* differential equation allowing to calculate the trajectory of a point charge interacting with an assigned “external” field $F_{\text{ext}}^{\mu\nu}$ (given in general by an incident radiation field plus the sum of the retarded fields generated by other charges). On the contrary, in our formulation of the theory Eq. (2.20) is a *second-order* equation which alone cannot determine the particle’s trajectory, since it is intrinsically coupled with the Eqs. (2.24) and (2.25) for the total field $F^{\mu\nu}$. The three equations need be solved together in order to determine the temporal evolution of the system of interacting particles and fields. One may observe incidentally that these equations are manifestly invariant under time reversal, which is not the case for the Lorentz-Dirac equation written in the form (2.31).

The solution of the system of coupled equations (2.20) and (2.24)–(2.25) depends on a set of initial conditions that is entirely specified by the values, at a given time $\bar{t}$, of $\mathbf{z}_n(\bar{t})$, $\mathbf{v}_n(\bar{t})$ and $F^{\mu\nu}(\mathbf{x}, \bar{t})$ for every $\mathbf{x}$. These initial conditions are subject to some relevant constraints, in addition to those already expressed by Eqs. (2.26) and (2.27). The reasonable requirement that $\bar{F}^{\mu\nu}(x)$ be non-singular at z_n implies that the divergent part of $F^{\mu\nu}(\mathbf{x}, \bar{t})$ must have the form given by Eqs. (2.8)–(2.10). Therefore the term of order $|\mathbf{x} - \mathbf{z}_n(\bar{t})|^{-2}$ is completely determined by $\mathbf{v}_n(\bar{t})$. Furthermore, the term of order $|\mathbf{x} - \mathbf{z}_n(\bar{t})|^{-1}$ is related to $\mathbf{v}_n(\bar{t})$ and to the zeroth order term $\tilde{F}^{\mu\nu}(\mathbf{z}_n(\bar{t}))$, since the vector $\dot{u}_n$ appearing in Eq. (2.10) must coincide with that resulting from Eq. (2.20).

Another subtler constraint on the initial conditions is related to the well-known problem of the “run-away” solutions of the Lorentz-Dirac equation [1–3]. We remark incidentally that such solutions do not violate the conservation law given by Eq. (2.23). In fact the unlimited increase towards $+\infty$ of the particle’s kinetic energy is compensated by a corresponding decrease towards $-\infty$ of the electromagnetic energy P_{em}^0 (which is not a definite positive quantity, according to our definition), and analogous statements are valid for the three spatial components of the momentum. These solutions, which are therefore completely legitimate from a mathematical standpoint, are generally dismissed for phenomenological reasons, since run-away electrons have never been observed.

We remind that, in the usual treatment of the Lorentz-Dirac equation as a third-order differential equation, the initial particle acceleration has to be specified among the initial conditions, together with the velocity, the position and the incident field. One may then take advantage of this additional free parameter in order to get rid of the run-away solutions. For instance, considering a scattering problem between a particle and a wave-packet initially very far apart, it is possible in general to obtain a finite result for the asymptotic momentum of the particle in the final state by introducing a suitable “preacceleration”, i.e. an acceleration of the particle that decreases as $\exp(t/\gamma\tau_0)$ for $t \rightarrow -\infty$, with $\tau_0 = q^2/6\pi mc^3$ ($\simeq 0.62 \times 10^{-23}$ sec in the case of an electron), and that becomes significant starting from a time of order τ_0 before the actual interaction between particle and wave-packet takes place.

In our formulation of the Cauchy problem for the equations (2.20) and (2.24)–(2.25), since the particle equation of motion is now of second order, the initial acceleration cannot be independently assigned, but is determined by the initial value of the total electromagnetic field according to Eq. (2.20). Therefore the requirement that the asymptotic particle momenta be finite both for $t \rightarrow -\infty$ and for $t \rightarrow +\infty$ is equivalent to a constraint on the initial values of the field. The fact that, in the traditional “third-order” treatment, the preacceleration is exponentially vanishing for times t very far in the past with respect to the actual interaction, within the new “second-order” picture translates into the fact that an equivalently small “adjustment” of an arbitrarily assigned field configuration at the same times t is generally required in order to satisfy the asymptotic conditions.

It is finally worth mentioning that the run-away behavior cannot be revealed by a perturbative expansion in the coupling constant. Therefore the existence of run-away solutions cannot be claimed to be a formal drawback of the classical theory of point charges as compared to perturbative quantum electrodynamics. It has also been conjectured that a particular kind of run-away solutions may exist in relativistic spin 1/2 QED [11].

III. THE ACTION PRINCIPLE

A. The renormalized lagrangian

A fundamental property of the system of differential equations constituted by Eqs. (2.20) and (2.24)–(2.27) is that it can be derived from an action principle. A lagrangian formalism can in fact be introduced in a natural way by taking as configuration variables of the system the position $\mathbf{z}_n(t)$ of the particles and the four-vector electromagnetic potential $A^\mu(\mathbf{x}, t)$ satisfying the relation

$$F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu. \quad (3.1)$$

Given an inertial system of coordinates, we shall define as the total lagrangian of the system at the time t the functional

$$L = L_{\text{em}} + L_{\text{part}} + L_{\text{int}}, \quad (3.2)$$

with

$$L_{\text{em}} = -\frac{1}{4}\mathcal{R} \int d^3\mathbf{x} F_{\mu\nu}(\mathbf{x}, t) F^{\mu\nu}(\mathbf{x}, t), \quad (3.3)$$

$$L_{\text{part}} = - \sum_{n=1}^N \frac{m_n}{\gamma_n}, \quad (3.4)$$

$$L_{\text{int}} = \sum_{n=1}^N \frac{q_n}{\gamma_n} u_n^\nu \tilde{A}_\nu(z_n). \quad (3.5)$$

The vector $\tilde{A}^\nu(z_n)$ in Eq. (3.5) is defined in perfect analogy with Eq. (2.18):

$$\tilde{A}^\nu(z_n) \equiv \mathcal{R} \lim_{r \rightarrow 0} \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} A^\nu(\mathbf{z}_n + \mathbf{x}, t). \quad (3.6)$$

One can see therefore that L is indeed a functional of the configuration variables and of their first time derivatives, taken at a given time t . In general this functional is defined for every field configuration having a behavior near the singular points such that the $\mathcal{R}$ -integral in Eq. (3.3) and the $\mathcal{R}$ -limit in Eq. (3.6) are properly defined.

As usual, Eqs. (2.24) and (2.27) are automatically satisfied as a result of Eq. (3.1), so that the equations of motion for the variables $\mathbf{z}_n$ and $A^\mu(x)$ reduce to Eqs. (2.20), (2.25) and (2.26). We are now going to show that these equations follow from Hamilton's variational principle.

Theorem 3 *If one defines the action I as*

$$I = \int_{t_1}^{t_2} L dt,$$

with L given by Eqs. (3.2)–(3.5), then the equations of motion of the system in the time interval $t_1 < t < t_2$ are equivalent to the condition that $\delta I = 0$ for every infinitesimal variation $\delta A^\mu(x)$ and $\delta \mathbf{z}_n(t)$ vanishing at $t = t_1$ and $t = t_2$, and such that the varied trajectories $\{\mathbf{z}_n + \delta \mathbf{z}_n, A^\mu + \delta A^\mu\}$ remain in the definition domain of I .

In order to prove this theorem, we shall introduce the non-renormalized lagrangian

$$\overline{L}(r) \equiv \overline{L}_{\text{em}}(r) + L_{\text{part}} + \overline{L}_{\text{int}}(r), \quad (3.7)$$

with

$$\overline{L}_{\text{em}}(r) = -\frac{1}{4} \int d^3\mathbf{x} F_{\mu\nu}(\mathbf{x}, t) F^{\mu\nu}(\mathbf{x}, t) \prod_{n=1}^N \theta(|\mathbf{x} - \mathbf{z}_n| - r) \quad (3.8)$$

and

$$\overline{L}_{\text{int}}(r) = \sum_{n=1}^N \frac{q_n}{\gamma_n} u_n^\nu \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} A_\nu(\mathbf{z}_n + \mathbf{x}, t). \quad (3.9)$$

Since $L = \mathcal{R} \lim_{r \rightarrow 0} \overline{L}(r)$, we can write

$$\overline{L}(r) = (L)_{-1} r^{-1} + L + O(r), \quad (3.10)$$

where the coefficient $(L)_{-1}$ is some other functional of the configuration variables and of their first time derivatives. It follows from Eq. (3.10) that

$$\delta \overline{L}(r) = \delta (L)_{-1} r^{-1} + \delta L + O(r). \quad (3.11)$$

Therefore the variations of the renormalized lagrangian L will be evaluated by calculating the corresponding variations of $\overline{L}$, and then taking their $\mathcal{R}$ -limit for $r \rightarrow 0$.

B. Derivation of the field equations from the action principle

Let us fix at a first stage $\delta \mathbf{z}_n \equiv 0$ and consider only variations of the electromagnetic potential, which will be represented by arbitrary infinitesimal test functions $\delta A^\mu(x)$ satisfying the boundary conditions. We want to show that, in order to have identically $\delta I = 0$, the fields $F^{\mu\nu}$ must define distributions that satisfy the differential equations (2.25) and (2.26). For $\delta \mathbf{z}_n \equiv 0$ we have

$$\delta \bar{L}(r) = - \int d^3 \mathbf{x} F^{\mu\nu} \partial_\mu \delta A_\nu + \sum_{n=1}^N \frac{q_n}{\gamma_n} u_n^\nu \delta A_\nu(z_n) + O(r), \quad (3.12)$$

where we have used the fact that, assuming $F^{\mu\nu}$ to have at most a singularity of order r^{-2} in z_n , the integral on the right-hand side is convergent, and moreover

$$\int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} \delta A_\nu(\mathbf{z}_n + \mathbf{x}, t) = \delta A_\nu(z_n) + O(r^2).$$

Since on the right-hand side of Eq. (3.12) there is no term of order r^{-1} , a comparison with Eq. (3.11) shows that $\delta(L)_{-1} = 0$ and

$$\delta I = \int_{t_1}^{t_2} dt \delta L = \int_{t_1}^{t_2} dt \left[- \int d^3 \mathbf{x} F^{\mu\nu} \partial_\mu \delta A_\nu + \sum_{n=1}^N \frac{q_n}{\gamma_n} u_n^\nu \delta A_\nu(z_n) \right]. \quad (3.13)$$

Since δA_ν are test functions vanishing for $t = t_1$ and $t = t_2$, Eq. (3.13) can be rewritten as

$$\delta I = \int_{t_1}^{t_2} dt \int d^3 \mathbf{x} \left[\partial_\mu F^{\mu\nu} + \sum_{n=1}^N \frac{q_n}{\gamma_n} u_n^\nu \delta(\mathbf{x} - \mathbf{z}_n) \right] \delta A_\nu,$$

where the partial derivatives of $F^{\mu\nu}$ are defined according to the distribution theory. Therefore the condition $\delta I = 0$ for arbitrary δA_ν implies

$$\partial_\mu F^{\mu\nu} = - \sum_{n=1}^N \frac{q_n}{\gamma_n} u_n^\nu \delta(\mathbf{x} - \mathbf{z}_n) = -j^\nu, \quad (3.14)$$

which is equivalent to Eqs. (2.25) and (2.26). Of course, assuming that the fields $F^{\mu\nu}$ are represented by ordinary differentiable functions of x everywhere outside the world-lines of the particles, Eq. (3.14) implies in particular that in the same region such functions satisfy the differential equation $\partial_\mu F^{\mu\nu} = 0$ in the usual sense.

It is well-known that the general solution of Eq. (3.14), when the particle trajectories are assigned, is given by the sum of the Liénard-Wiechert fields and a solution of the free-field equations. Using this fact, we are now going to show that the lagrangian L is indeed correctly defined on the solutions of the field equations and on the neighboring trajectories obtained via non-singular variations δA_ν of the potential. This fact ensures the self-consistency of the whole procedure. First of all, since asymptotic conditions of the type (2.8) must be satisfied by the fields near the particles, we can repeat the same considerations we made about the $\mathcal{R}$ -integral in Eq. (2.6), showing that also the right-hand side of Eq. (3.3) is correctly defined. Moreover, using the expansion for the advanced and retarded vector potentials derived in

App. A, one can verify that also the potential $\tilde{A}_\nu(z_n)$ appearing in Eq. (3.5) is well defined according to Eq. (3.6). In particular, defining for a given particle

$$\overline{A}^\mu \equiv A^\mu - \frac{1}{2}(A_{\text{adv}}^\mu + A_{\text{ret}}^\mu), \quad (3.15)$$

one gets from Eqs. (A6) and (A8)

$$A^\mu(z+x) = (A^\mu)_{-1}(x) + (A^\mu)_0(x) + \overline{A}^\mu(z) + O(x), \quad (3.16)$$

and from the antisymmetry in x of $(A^\mu)_0(x)$ it follows that

$$\tilde{A}^\mu(z) = \lim_{r \rightarrow 0} \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} [(A^\mu)_0(x) + \overline{A}^\mu(z)] = \overline{A}^\mu(z), \quad (3.17)$$

in close analogy with Eqs. (2.28)–(2.30). We remark also that Eqs. (3.15) and (3.17) ensure that $\tilde{A}^\mu(z)$ behaves as a four-vector under Lorentz transformations.

C. Derivation of the particle equation of motion from the action principle

When considering also variations $\delta \mathbf{z}_n(t) \neq 0$, one has to pay attention to the fact that the quantity $\tilde{A}_\nu(z_n)$ is defined only on the world-line of a particle, so that its partial derivatives with respect to $\mathbf{z}_n$ are not defined in the usual sense. This is equivalent to the obvious statement that, whenever $\delta \mathbf{z}_n(t) \neq 0$ and $\delta A^\mu(x) = 0$, the varied trajectories no longer belong to the definition domain of the functional L . In order to remain within this domain, each variation $\delta \mathbf{z}_n(t) \neq 0$ must be accompanied by a simultaneous variation δA^μ of the potentials, characterized by an appropriate singular behavior near the particles. For this reason, it is convenient to proceed in two steps. First, given the boundary conditions at $t = t_1$ and $t = t_2$, for each possible trajectory z_n of the particles one determines the potential $A_{z_n}^\mu(x)$ that minimizes the action I with respect to the variations of the potential alone: as we have seen in the previous subsection, $A_{z_n}^\mu(x)$ is just the potential that solves the Maxwell equations for those particular particle trajectories. Then one can define the functional $I'[z_n]$ of the trajectory of the particles alone, as

$$I'[z_n] \equiv I[z_n, A_{z_n}^\mu(x)]. \quad (3.18)$$

It is now possible to show that the condition $\delta I' = 0$, for arbitrary variations $\delta \mathbf{z}_n(t)$ satisfying $\delta \mathbf{z}_n(t_1) = \delta \mathbf{z}_n(t_2) = 0$, is equivalent to the fact that $\mathbf{z}_n(t)$, together with $A_{z_n}^\mu(x)$, are solutions of Eq. (2.20). Of course, the minima of I must coincide with the minima of I' : therefore the trajectory of particles and fields $\{z_n, A_{z_n}^\mu(x)\}$ satisfying Eq. (2.20) is indeed the trajectory that minimizes I within its whole definition domain, subject to the assigned boundary conditions.

Keeping these ideas in mind, we shall still write for simplicity A^μ instead of $A_{z_n}^\mu$, although we shall always implicitly assume that the field satisfies Eq. (3.14). We shall therefore apply the previous formulae describing the asymptotic behavior of the field near the point charge. We shall make use also of an expansion for the derivatives of the potentials that, similarly to Eq. (2.29), follows from the results of App. A:

$$\partial_\mu A_\nu(z+x) = (\partial_\mu A_\nu)_{-2}(x) + (\partial_\mu A_\nu)_{-1}(x) + (\partial_\mu A_\nu)_0(x) + \partial_\mu \tilde{A}_\nu(z) + O(x), \quad (3.19)$$

where $(\partial_\mu A_\nu)_j(x)$ is a homogeneous function of degree j and even (resp. odd) parity for j odd (resp. even), while

$$\partial_\mu \tilde{A}_\nu(z) \equiv \mathcal{R} \lim_{r \rightarrow 0} \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} \partial_\mu A_\nu(\mathbf{z} + \mathbf{x}, t). \quad (3.20)$$

Note that the above equation is to be considered as a definition since, as we already observed, the partial derivatives of $\tilde{A}(z)$ do not exist in the usual sense. However one can easily show that, in close analogy with Eqs. (2.30) and (3.17),

$$\partial_\mu \tilde{A}_\nu(z) = \partial_\mu \overline{A}_\nu(z), \quad (3.21)$$

where the right-hand side represents the usual partial derivative of the function $\overline{A}(x)$ defined by Eq. (3.15).

With a rather tedious but straightforward calculation, which is reported in App. E, one obtains

$$\begin{aligned} \delta L &= -\frac{d}{dx^0} \left(\mathcal{R} \int d^3\mathbf{x} F^{0j} \delta A_j \right) \\ &\quad + \sum_{n=1}^N \left[q_n \tilde{A}_i(z_n) \delta v_n^i + q_n \gamma_n^{-1} u_n^\nu \partial_i \tilde{A}_\nu(z_n) \delta z_n^i + m_i u_{n,i} \delta v_n^i \right] \\ &= \frac{d}{dt} \left[\sum_{n=1}^N \left(m_n u_{n,i} + q_n \tilde{A}_i(z_n) \right) \delta z_n^i - \mathcal{R} \int d^3\mathbf{x} F^{0j} \delta A_j \right] \\ &\quad - \sum_{n=1}^N \left[\frac{d}{dt} \left(m_n u_{n,i} + q_n \tilde{A}_i(z_n) \right) - q_n \gamma_n^{-1} u_n^\nu \partial_i \tilde{A}_\nu(z_n) \right] \delta z_n^i. \end{aligned} \quad (3.22)$$

Since $\delta z_n^i = 0$ and $\delta A_\nu = 0$ for $t = t_1$ and $t = t_2$, the boundary terms do not contribute to the variation of the action, and we obtain

$$\delta I = - \int_{t_1}^{t_2} dt \sum_{n=1}^N \left[\frac{d}{dt} \left(m_n u_{n,i} + q_n \tilde{A}_i(z_n) \right) - q_n \gamma_n^{-1} u_n^\nu \partial_i \tilde{A}_\nu(z_n) \right] \delta z_n^i. \quad (3.23)$$

Since it follows from Eqs. (3.17) and (3.21) that

$$\frac{d}{dt} \tilde{A}_\mu(z_n) = \frac{d}{dt} \overline{A}_\mu(z_n) = \gamma_n^{-1} u_n^\nu \partial_\nu \overline{A}_\mu(z_n) = \gamma_n^{-1} u_n^\nu \partial_\nu \tilde{A}_\mu(z_n),$$

the condition $\delta I = 0$ for arbitrary δz_n^i implies

$$m_n \dot{u}_{n,i} = m_n \gamma_n \frac{du_{n,i}}{dt} = q_n \left(\partial_i \tilde{A}_\nu(z_n) - \partial_\nu \tilde{A}_i(z_n) \right) u_n^\nu = q_n \tilde{F}_{i\nu}(z_n) u_n^\nu, \quad (3.24)$$

that coincides with Eq. (2.20). We have therefore proved that the dynamics of the system is entirely determined by Hamilton's variational principle applied to the lagrangian defined by Eqs. (3.2)–(3.5), as stated by Theorem 3. Of course, it is also possible to deduce directly the conservation laws of energy and momentum from the translational invariance in space and time of L . This is illustrated in detail in App. F.

D. A comparison with the idea of mass renormalization

The renormalization of our lagrangian was achieved simply through the introduction of a $\mathcal{R}$ -integral for the electromagnetic part in Eq. (3.3) and a $\mathcal{R}$ -limit of the electromagnetic potential for the interaction part in Eq. (3.5). Clearly this procedure has in principle nothing to do with the renormalization of the particle's mass, which is typical of other approaches to the argument, and in particular of standard renormalization techniques of quantum field theories. Therefore no infinitely negative “bare mass” has to be introduced in our theory, and the difficulty of simultaneously compensating with such a mass term the non-covariant divergent parts in Eqs. (2.15)–(2.16) is automatically avoided. It is however interesting to note that the divergent part of Eq. (3.8) can be easily calculated using Eq. (B9):

$$\begin{aligned} (L_{\text{em}})_{-1} r^{-1} &= -\frac{1}{4} \int d^3\mathbf{x} (F^{\mu\nu})_{-2} (F_{\mu\nu})_{-2} \theta(y-r) \\ &= -\frac{1}{4\gamma} \int_r^\infty y^2 dy \int d\Omega_{\mathbf{y}} \left[-\frac{2q^2}{(4\pi)^2 y^4} \right] = \frac{q^2}{8\pi\gamma r} \end{aligned}$$

(we consider here for simplicity a single charged particle), so that we have

$$L_{\text{em}} = \lim_{r \rightarrow 0} \left[-\frac{1}{4} \int d^3\mathbf{x} F^{\mu\nu} F_{\mu\nu} \theta(y-r) - \frac{q^2}{8\pi\gamma r} \right]. \quad (3.25)$$

Similarly we have from Eq. (A6)

$$\int_{y=r} d\Omega_{\mathbf{y}} (A_\nu)_{-1} = \int_{y=r} d\Omega_{\mathbf{y}} \frac{qu_\nu}{4\pi y} = \frac{qu_\nu}{r},$$

so that

$$L_{\text{int}} = \frac{q}{\gamma} u^\nu \mathcal{R} \lim_{r \rightarrow 0} \int_{y=r} \frac{d\Omega_{\mathbf{y}}}{4\pi} A_\nu(\mathbf{z} + \mathbf{x}) = \lim_{r \rightarrow 0} \left[\frac{q}{\gamma} u^\nu \int_{y=r} \frac{d\Omega_{\mathbf{y}}}{4\pi} A_\nu(\mathbf{z} + \mathbf{x}) + \frac{q^2}{4\pi r \gamma} \right]. \quad (3.26)$$

Therefore, combining Eqs. (3.2), (3.25) and (3.26), we can write

$$L = \lim_{r \rightarrow 0} \left[-\frac{1}{4} \int d^3\mathbf{x} F^{\mu\nu} F_{\mu\nu} \theta(y-r) + \frac{q}{\gamma} u^\nu \int_{y=r} \frac{d\Omega_{\mathbf{y}}}{4\pi} A_\nu(\mathbf{z} + \mathbf{x}) - \frac{m_0}{\gamma} \right], \quad (3.27)$$

with

$$m_0 = m - \frac{q^2}{8\pi r}.$$

Clearly, one could interpret Eq. (3.27) as the lagrangian of a particle having indeed a divergent negative bare mass m_0 and carrying a total electric charge q that, for an observer with the same instantaneous velocity of the particle, looks uniformly distributed over an infinitesimal spherical surface of radius r . We must however underline that Eq. (3.27) is verified only when the fields are already solutions of Maxwell equations. In general, taking Eq. (3.27) as a definition would not be equivalent to the definition of L that we have given in Eqs. (3.2)–(3.5). This becomes particularly evident when introducing the hamiltonian formalism. In fact, since each of the three terms on the right-hand side of Eq. (3.27) is divergent, it is immediate to see that also the conjugate momenta of the particles $p_i = \partial L / \partial v^i$ are divergent in the $r \rightarrow 0$ limit. On the contrary, the lagrangian (3.2) is the sum of three *individually finite* terms. As we shall show in the next section, from this fact it follows that all the momenta are finite, and therefore an exact hamiltonian can be defined for the theory.

IV. EULER-LAGRANGE AND HAMILTON EQUATIONS

It is useful to define the partial derivatives of the lagrangian L with respect to its arguments, in such a way that the equations of motion can be formally derived as the usual Euler-Lagrange equations associated with L . We have seen in the previous section that, in order to remain within the original definition domain (that we shall call D_0) of L , each variation δz_n must be accompanied by a variation δA of the electromagnetic potential with an appropriate singular behavior near the particles. Therefore, defining a partial derivative of the type $\partial L / \partial z_n^i$ for constant potential $A^\mu(x)$ requires introducing an extension of L outside D_0 . We have to note first of all that, inside D_0 , the particles' positions z_n are univocally determined by the field variables $A^\mu(x)$, since the potential is assumed to be singular just in z_n . In other words, as soon as $A^\mu(\mathbf{x}, t)$ is assigned for every $\mathbf{x}$, also its singular points, that will be now called $\mathbf{w}_n(t)$ to distinguish them in general from the particles' positions $\mathbf{z}_n(t)$, are of course determined, and inside D_0 it is always assumed that $\mathbf{z}_n = \mathbf{w}_n$. It is clear now that, in order to treat $\mathbf{z}_n$ and $A^\mu(x)$ as really independent variables, we have to define L also for $\mathbf{z}_n \neq \mathbf{w}_n$. Having this in mind, the results of the previous section indicate how to extend in an obvious way the definition of L . We shall put

$$\frac{\partial L_{\text{em}}}{\partial z_n^i} = 0, \quad (4.1)$$

which is consistent with considering L_{em} as the purely electromagnetic part of the lagrangian, and

$$\frac{\partial L_{\text{int}}}{\partial z_n^i} = q_n \frac{u_n^\nu}{\gamma_n} \partial_i \tilde{A}_\nu(z_n), \quad (4.2)$$

which is equivalent to extending in a neighborhood of $\mathbf{w}_n$ the function $\tilde{A}_\nu(z_n)$ appearing in Eq. (3.5) according to

$$\tilde{A}_\nu(\mathbf{z}_n, t) = \tilde{A}_\nu(\mathbf{w}_n, t) + \partial_i \tilde{A}_\nu(\mathbf{w}_n, t)(z_n^i - w_n^i) + O[(\mathbf{z}_n - \mathbf{w}_n)^2].$$

On the right-hand side of the above equation the quantities $\tilde{A}_\nu(\mathbf{w}_n, t)$ and $\partial_i \tilde{A}_\nu(\mathbf{w}_n, t)$ are defined according to Eqs. (3.6) and (3.20) respectively, whereas terms of order $(\mathbf{z}_n - \mathbf{w}_n)^2$ can be neglected for our purposes, since they clearly do not affect the equations of motion. Besides, the derivatives with respect to $v_n^i = dz_n^i/dt$ can be determined unambiguously as

$$\begin{aligned} \frac{\partial L_{\text{em}}}{\partial v_n^i} &= 0, \\ \frac{\partial L_{\text{int}}}{\partial v_n^i} &= q \tilde{A}_i(z_n). \end{aligned}$$

We have then, as a result of Eq. (3.24),

$$\frac{\partial L}{\partial z_n^i} = q_n \frac{u_n^\nu}{\gamma_n} \partial_i \tilde{A}_\nu(z_n) = \frac{d}{dt} [m_n u_{n,i} + q_n \tilde{A}_i(z_n)] = \frac{d}{dt} \frac{\partial L}{\partial v_n^i}.$$

Similarly, using Eq. (3.14), we have

$$\frac{\delta L}{\delta A_\mu(x)} = \partial_j F^{j\mu}(x) + \sum_{n=1}^N q_n \frac{u_n^\mu}{\gamma_n} \delta(\mathbf{x} - \mathbf{z}_n) = -\partial_0 F^{0\mu}(x) = \frac{d}{dt} \frac{\delta L}{\delta \partial_0 A_\mu(x)},$$

where the functional derivatives of L with respect to $A_\mu(x)$ and $\partial_0 A_\mu(x)$ are unambiguously defined as distributions in the usual way. Looking at the first and last members of the two above equations, the formal structure of Euler-Lagrange equations is directly recognizable.

The hamiltonian functional can be introduced via a Legendre transformation on L , as in ordinary analytical mechanics. As usual, after imposing the Lorentz gauge condition

$$\partial_\mu A^\mu(x) = 0,$$

we shall add a term $-(1/2) (\partial_\mu A^\mu)^2$ to the electromagnetic lagrangian density in order to introduce a conjugate momentum also for the temporal component $A^0(x)$ of the potential. We shall therefore replace Eq. (3.3) with

$$L_{\text{em}} = -\mathcal{R} \int d^3\mathbf{x} \left\{ \frac{1}{4} F_{\mu\nu}(\mathbf{x}, t) F^{\mu\nu}(\mathbf{x}, t) + \frac{1}{2} [\partial_\mu A^\mu(\mathbf{x}, t)]^2 \right\}.$$

As a result, the Euler-Lagrange equation for A^μ becomes

$$\partial_\nu \partial^\nu A^\mu(x) = - \sum_{n=1}^N q_n \frac{u_n^\mu}{\gamma_n} \delta(\mathbf{x} - \mathbf{z}_n) = -j^\mu(x). \quad (4.3)$$

Then, after introducing the canonical momenta

$$p_{n,i} \equiv \frac{\partial L}{\partial v_n^i} = m_n u_{n,i} + q_n \tilde{A}_i(z_n) \quad (4.4)$$

$$\pi_i(x) \equiv \frac{\delta L}{\delta \partial_0 A^i(x)} = F_{0i}(x) \quad (4.5)$$

$$\pi_0(x) \equiv \frac{\delta L}{\delta \partial_0 A^0(x)} = \partial_\mu A^\mu(x), \quad (4.6)$$

we can define the hamiltonian H of the system as

$$\begin{aligned} H &\equiv \sum_{n=1}^N p_{n,i} v_n^i + \mathcal{R} \int d^3\mathbf{x} \pi_\mu \partial_0 A^\mu - L \\ &= \mathcal{R} \int d^3\mathbf{x} \left(\frac{1}{4} F_{ij} F^{ij} + \frac{1}{2} \pi_\mu \pi^\mu - \pi^i \partial_i A^0 + \pi^0 \partial_i A^i \right) \\ &\quad + \sum_{n=1}^N \left[q_n \tilde{A}^0(z_n) + \sqrt{(p_{n,i} - q_n \tilde{A}_i(z_n)) (p_n^i - q_n \tilde{A}^i(z_n)) + m_n^2} \right]. \end{aligned} \quad (4.7)$$

Hamilton's equations are:

$$\frac{dz_n^i}{dt} = \frac{\partial H}{\partial p_{n,i}} = \frac{p_n^i - q_n \tilde{A}^i(z_n)}{\sqrt{(p_{n,k} - q_n \tilde{A}_k(z_n)) (p_n^k - q_n \tilde{A}^k(z_n)) + m_n^2}}, \quad (4.8)$$

$$\partial_0 A^i(x) = \frac{\delta H}{\delta \pi_i(x)} = \pi^i(x) + \partial^i A_0(x), \quad (4.9)$$

$$\partial_0 A^0(x) = \frac{\delta H}{\delta \pi_0(x)} = \pi^0(x) - \partial_i A^i(x), \quad (4.10)$$

$$\frac{dp_n^i}{dt} = -\frac{\partial H}{\partial z_{n,i}} = \frac{q_n \partial^i \tilde{A}^j(z_n) (p_{n,j} - q_n \tilde{A}_j(z_n))}{\sqrt{(p_{n,k} - q_n \tilde{A}_k(z_n)) (p_n^k - q_n \tilde{A}^k(z_n)) + m_n^2}} - q_n \partial^i \tilde{A}^0(z_n), \quad (4.11)$$

$$\begin{aligned} \partial_0 \pi^i(x) = -\frac{\delta H}{\delta A_i(x)} = \partial_j F^{ji}(x) + \partial^i \pi^0(x) \\ + \sum_{n=1}^N \frac{q_n (p_n^i - q_n \tilde{A}^i(z_n)) \delta(\mathbf{x} - \mathbf{z}_n)}{\sqrt{(p_{n,k} - q_n \tilde{A}_k(z_n)) (p_n^k - q_n \tilde{A}^k(z_n)) + m_n^2}}, \end{aligned} \quad (4.12)$$

$$\partial_0 \pi^0(x) = -\frac{\delta H}{\delta A_0(x)} = \partial_i \pi^i(x) + \sum_{n=1}^N q_n \delta(\mathbf{x} - \mathbf{z}_n). \quad (4.13)$$

As usual, Eqs. (4.8)–(4.10) are just the inversion of Eqs. (4.4)–(4.6), whereas Eq. (4.11) is equivalent to Eq. (3.24) and Eqs. (4.12)–(4.13) are equivalent to Eq. (4.3).

The theory can be expressed by means of fundamental Poisson brackets between canonical variables having the familiar form

$$\{z_n^i, p_m^j\} = \delta_{nm} \delta^{ij} \quad (4.14)$$

$$\{A^\mu(\mathbf{x}), \pi^\nu(\mathbf{y})\} = g^{\mu\nu} \delta(\mathbf{x} - \mathbf{y}), \quad (4.15)$$

all the others being equal to zero. After introducing the total momentum

$$P^i \equiv -\mathcal{R} \int d^3\mathbf{x} \pi_\mu \partial^i A^\mu + \sum_{n=1}^N p_n^i \quad (4.16)$$

and putting $P^0 \equiv H$, it is straightforward to verify that from Eqs. (4.8)–(4.15) it follows

$$\begin{aligned} \{z_n^i, P^0\} &= \frac{\partial P^0}{\partial p_{n,i}} = \frac{dz_n^i}{dt} \\ \{z_n^i, P^j\} &= \frac{\partial P^j}{\partial p_{n,i}} = \delta^{ij} \\ \{p_n^i, P^0\} &= -\frac{\partial P^0}{\partial z_{n,i}} = \frac{dp_n^i}{dt} \\ \{p_n^i, P^j\} &= -\frac{\partial P^j}{\partial z_{n,i}} = 0 \\ \{A^\mu(x), P^\nu\} &= \frac{\delta P^\nu}{\delta \pi_\mu(x)} = -\partial^\nu A^\mu(x) \\ \{\pi^\mu(x), P^\nu\} &= -\frac{\delta P^\nu}{\delta A_\mu(x)} = -\partial^\nu \pi^\mu(x), \end{aligned}$$

where the members of the last two equations are to be interpreted as distributions. The above equations show that, as expected, P^μ is the generator of the space-time translations for the system. Of course, on the solutions of the equations of motion it is equal to the total energy-momentum introduced in Sec. II. This is explicitly shown in App. F.

V. THE RENORMALIZATION PROCEDURE IN THE WAVE-VECTOR SPACE

A. The energy-momentum in terms of the Fourier transform of the electromagnetic field

Let us operate the Fourier decomposition of the electromagnetic potential according to the formulae

$$A_\mu(x) = \frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} \left[a_\mu(\mathbf{k}, t) e^{i\mathbf{k}\cdot\mathbf{x}} + a_\mu^*(\mathbf{k}, t) e^{-i\mathbf{k}\cdot\mathbf{x}} \right] \quad (5.1a)$$

$$\partial_0 A_\mu(x) = -\frac{i}{2(2\pi)^3} \int d^3\mathbf{k} \left[a_\mu(\mathbf{k}, t) e^{i\mathbf{k}\cdot\mathbf{x}} - a_\mu^*(\mathbf{k}, t) e^{-i\mathbf{k}\cdot\mathbf{x}} \right], \quad (5.1b)$$

with $k^0 \equiv |\mathbf{k}|$. As a consequence of Eqs. (4.15) and (5.1), the Poisson brackets between the variables a_μ and a_μ^* are

$$\{a_\mu(\mathbf{k}), a_\nu(\mathbf{k}')\} = \{a_\mu^*(\mathbf{k}), a_\nu^*(\mathbf{k}')\} = 0 \quad (5.2a)$$

$$\{a_\mu(\mathbf{k}), a_\nu^*(\mathbf{k}')\} = -2(2\pi)^3 i g_{\mu\nu} k^0 \delta(\mathbf{k} - \mathbf{k}'). \quad (5.2b)$$

It is well-known that for a free field one can express the electromagnetic energy-momentum as

$$P_{\text{free}}^\mu = \frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} k^\mu a^\nu(\mathbf{k}) a_\nu^*(\mathbf{k}).$$

It is not surprising that, when point-like charged particles are present, also the above integral becomes divergent. The same is true for the integrals on the right-hand side of the Fourier expansions (5.1) when $\mathbf{x}$ takes the value of the position $\mathbf{z}_n(t)$ of one of the particles. This can be verified using the following asymptotic expansion of $a^\nu(\mathbf{k}, t)$ for $|\mathbf{k}| \rightarrow \infty$, which is derived in App. G:

$$a^\nu(\mathbf{k}, t) = \sum_{n=1}^N \left[(a_n^\nu)_{-1}(\mathbf{k}, t) + (a_n^\nu)_0(\mathbf{k}, t) \right] e^{-i\mathbf{k}\cdot\mathbf{z}_n} + O(|\mathbf{k}|^{-3}), \quad (5.3)$$

with

$$(a_n)_{-1} = -q_n \frac{u_n}{k \cdot u_n}, \quad (5.4)$$

$$(a_n)_0 = -iq_n \frac{(k \cdot \dot{u}_n) u_n - (k \cdot u_n) \dot{u}_n}{(k \cdot u_n)^3}. \quad (5.5)$$

Clearly, Eq. (5.3) is the direct counterpart of the expansions (3.16) and (3.19) of $A^\nu(z+x)$ and $\partial_\mu A^\nu(z+x)$ for $|\mathbf{x}| \rightarrow 0$. We have in particular

$$(A_n^\mu)_j(x) = \frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} \left[(a_n^\mu)_j(\mathbf{k}, t) e^{i\mathbf{k}\cdot\mathbf{x}} + (a_n^\mu)_j^*(\mathbf{k}, t) e^{-i\mathbf{k}\cdot\mathbf{x}} \right] \quad (5.6)$$

$$(\partial_\nu A_n^\mu)_{j-1}(x) = \frac{i}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} k_\nu \left[(a_n^\mu)_j(\mathbf{k}, t) e^{i\mathbf{k}\cdot\mathbf{x}} - (a_n^\mu)_j^*(\mathbf{k}, t) e^{-i\mathbf{k}\cdot\mathbf{x}} \right]. \quad (5.7)$$

It can be noticed that $(a_n)_j$ is a homogeneous function of $\mathbf{k}$ of degree $-j-2$. Besides, it is real for $j = -1$ and purely imaginary for $j = 0$.

We can however redefine the divergent integrals in $\mathbf{k}$ space by means of a renormalization procedure equivalent to that introduced in Sec. II for the corresponding expressions in spatial coordinates.

Definition 2 *Given a function $f(\Lambda)$ having for $\Lambda \rightarrow \infty$ a behavior of the form*

$$f(\Lambda) = c\Lambda + d + O(\Lambda^{-1}),$$

we define its $\mathcal{R}$ -limit for $\Lambda \rightarrow \infty$ as

$$\mathcal{R} \lim_{\Lambda \rightarrow \infty} f(\Lambda) \equiv d.$$

Correspondingly, given a function $g(\mathbf{k})$ having a behavior such that

$$\int_{|\mathbf{k}| < \Lambda} d^3\mathbf{k} g(\mathbf{k}) = c'\Lambda + d' + O(\Lambda^{-1}) \quad \text{for } \Lambda \rightarrow \infty,$$

we define its $\mathcal{R}$ -integral over the $\mathbf{k}$ space as

$$\mathcal{R} \int d^3\mathbf{k} g(\mathbf{k}) \equiv \mathcal{R} \lim_{\Lambda \rightarrow \infty} \int_{|\mathbf{k}| < \Lambda} d^3\mathbf{k} g(\mathbf{k}) = d'. \quad (5.8)$$

Using this mathematical tool one can express in terms of the variables a_μ and a_μ^* the canonical energy-momentum four-vector for the system of interacting particles and field, as stated by the following theorem.

Theorem 4 *The four-vectors P^μ and $\tilde{A}^\mu(z_n)$, given respectively by Eqs. (4.7), (4.16) and (3.6), can be equivalently expressed by the equations*

$$P^i = \frac{1}{2(2\pi)^3} \mathcal{R} \int \frac{d^3\mathbf{k}}{k^0} k^i a^\nu(\mathbf{k}, t) a_\nu^*(\mathbf{k}, t) + \sum_{n=1}^N p_n^i, \quad (5.9)$$

$$\begin{aligned} P^0 = & \frac{1}{2(2\pi)^3} \mathcal{R} \int d^3\mathbf{k} a^\nu(\mathbf{k}, t) a_\nu^*(\mathbf{k}, t) \\ & + \sum_{n=1}^N \left[q_n \tilde{A}^0(z_n) + \sqrt{(p_{n,i} - q_n \tilde{A}_i(z_n)) (p_n^i - q_n \tilde{A}^i(z_n)) + m_n^2} \right] \end{aligned} \quad (5.10)$$

and

$$\tilde{A}_\mu(z_n) = \frac{1}{2(2\pi)^3} \mathcal{R} \int \frac{d^3\mathbf{k}}{k^0} \left[a_\mu(\mathbf{k}, t) e^{i\mathbf{k} \cdot \mathbf{z}_n} + a_\mu^*(\mathbf{k}, t) e^{-i\mathbf{k} \cdot \mathbf{z}_n} \right]. \quad (5.11)$$

The proof of Theorem 4 is given in App. H. By making use of Eqs. (5.2), (5.10) and (5.11), one readily obtains the equation of motion of the a^μ :

$$\begin{aligned}\frac{d}{dt}a^\mu(\mathbf{k}, t) &= \{a^\mu(\mathbf{k}, t), P^0\} = -ik^0 a^\mu(\mathbf{k}, t) + \frac{i}{2(2\pi)^3 k^0} \sum_{n=1}^N q_n \frac{u_n^\mu}{\gamma_n} e^{-i\mathbf{k}\cdot\mathbf{z}_n} \\ &= -ik^0 a^\mu(\mathbf{k}, t) + \frac{i}{2(2\pi)^3 k^0} \int d^3\mathbf{x} j^\mu(\mathbf{x}, t) e^{-i\mathbf{k}\cdot\mathbf{x}},\end{aligned}$$

which is of course in accordance with Eqs. (4.3) and (5.1). We remark also that, with exactly the same procedure followed to derive Eq. (5.11), one can prove that the quantity $\partial_\nu \tilde{A}_\mu(z_n)$ defined by Eq. (3.20) can be expressed as

$$\partial_\nu \tilde{A}_\mu(z_n) = \frac{i}{2(2\pi)^3} \mathcal{R} \int \frac{d^3\mathbf{k}}{k^0} k_\nu \left[a_\mu(\mathbf{k}, t) e^{i\mathbf{k}\cdot\mathbf{z}_n} - a_\mu^*(\mathbf{k}, t) e^{-i\mathbf{k}\cdot\mathbf{z}_n} \right].$$

B. Another remark on the renormalization of mass

We saw in Sec. II that the expression for the electromagnetic energy-momentum in $\mathbf{x}$ space can be most conveniently handled when the cut-off regions in the integration domain are taken as ellipsoids like in Eqs. (2.11) and (2.12). Similarly, in some cases the $\mathcal{R}$ -integrals in $\mathbf{k}$ space appearing in Eqs. (5.9)–(5.10) may be more suitably expressed by choosing a different cut-off instead of the sphere of radius Λ introduced in Eq. (5.8). A particularly interesting situation occurs when a single charged particle is present. In such a case, putting

$$\bar{a}^\nu(k, t) \equiv a^\nu(\mathbf{k}, t) e^{i\mathbf{k}\cdot\mathbf{z}},$$

with $k^0 \equiv |\mathbf{k}|$, it can be easily seen that $\bar{a}^\nu$ transforms as a four-vector, as it can already be argued from the form of the right-hand sides of Eqs. (5.4) and (5.5). It follows that also the expression

$$\bar{K}_{\text{em}}^\mu(t, \Lambda) \equiv \frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} k^\mu a^\nu(\mathbf{k}, t) a_\nu^*(\mathbf{k}, t) \theta(\Lambda + k_\lambda u^\lambda(t)) \quad (5.12)$$

transforms as a four-vector, since it can be rewritten as

$$\bar{K}_{\text{em}}^\mu(t, \Lambda) \equiv \frac{1}{(2\pi)^3} \int d^4k k^\mu \bar{a}^\nu(k, t) \bar{a}_\nu^*(k, t) \delta(k^2) \theta(k^0) \theta(\Lambda + k_\lambda u^\lambda(t)).$$

We observe that the integration domain in Eq. (5.12) is characterized by the condition

$$|\mathbf{k}| - \mathbf{k} \cdot \mathbf{v} < \frac{\Lambda}{\gamma},$$

which corresponds to the volume contained into a rotational ellipsoid with a focus at the origin. Repeating the procedure used in App. H to obtain Eqs. (H10) and (H11), it is easy to see that

$$\mathcal{R} \lim_{\Lambda \rightarrow \infty} \bar{K}_{\text{em}}^\mu(t, \Lambda) = \frac{1}{2(2\pi)^3} \mathcal{R} \int \frac{d^3\mathbf{k}}{k^0} k^\mu a^\nu(\mathbf{k}, t) a_\nu^*(\mathbf{k}, t).$$

Furthermore, for $\mathbf{v}(t) = 0$ we get from Eq. (5.4)

$$\begin{aligned} & \frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} k^\mu (a^\nu)_{-1} (a_\nu)^*_{-1} \theta(\Lambda + k_\lambda u^\lambda(t)) \\ &= -\frac{q^2}{2(2\pi)^3} \int d^3\mathbf{k} \frac{k^\mu}{(k^0)^3} \theta(\Lambda - k^0) = -\frac{q^2}{4\pi^2} \Lambda u^\mu(t). \end{aligned}$$

Thanks to the Lorentz covariance, we can immediately generalize the above result for arbitrary $\mathbf{v}(t)$, obtaining

$$\overline{K}_{\text{em}}^\mu(t, \Lambda) = -\frac{q^2}{4\pi^2} \Lambda u^\mu(t) + \frac{1}{2(2\pi)^3} \mathcal{R} \int \frac{d^3\mathbf{k}}{k^0} k^\mu a^\nu(\mathbf{k}, t) a_\nu^*(\mathbf{k}, t) + O(\Lambda^{-1}). \quad (5.13)$$

A similar procedure can be followed for the $\mathcal{R}$ -integral that expresses $\tilde{A}_\mu(z)$ according to Eq. (5.11). Applying again the covariant cut-off we can define the four-vector

$$\begin{aligned} A_\mu(z, \Lambda) &\equiv \frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} [a_\mu(\mathbf{k}, t) e^{i\mathbf{k}\cdot\mathbf{z}} + a_\mu^*(\mathbf{k}, t) e^{-i\mathbf{k}\cdot\mathbf{z}}] \theta(\Lambda + k_\lambda u^\lambda(t)) \\ &= \frac{1}{(2\pi)^3} \int d^4k [\bar{a}_\mu(k, t) + \bar{a}_\mu^*(k, t)] \delta(k^2) \theta(k^0) \theta(\Lambda + k_\lambda u^\lambda(t)). \end{aligned} \quad (5.14)$$

Then, proceeding as in Eqs. (H17)–(H18), it is immediate to see that

$$\tilde{A}_\mu(z) = \mathcal{R} \lim_{\Lambda \rightarrow \infty} A_\mu(z, \Lambda).$$

Moreover, since for $\mathbf{v}(t) = 0$

$$\begin{aligned} & \frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} [(a_\mu)_{-1}(\mathbf{k}, t) e^{i\mathbf{k}\cdot\mathbf{z}} + (a_\mu)^*_{-1}(\mathbf{k}, t) e^{-i\mathbf{k}\cdot\mathbf{z}}] \theta(\Lambda + k_\lambda u^\lambda(t)) \\ &= \frac{q}{(2\pi)^3} \int d^3\mathbf{k} \frac{u_\mu(t)}{(k^0)^2} \theta(\Lambda - k^0) = \frac{q}{2\pi^2} \Lambda u_\mu(t), \end{aligned}$$

the Lorentz covariance implies that for arbitrary $\mathbf{v}(t)$

$$A_\mu(z, \Lambda) = \frac{q}{2\pi^2} \Lambda u_\mu(t) + \tilde{A}_\mu(z) + O(\Lambda^{-1}) \quad \text{for } \Lambda \rightarrow \infty. \quad (5.15)$$

Putting together Eqs. (5.9), (5.10), (5.13) and (5.15), we can finally write

$$\begin{aligned} P^\mu &= m u^\mu(t) + q \tilde{A}^\mu(z) + \frac{1}{2(2\pi)^3} \mathcal{R} \int \frac{d^3\mathbf{k}}{k^0} k^\mu a^\nu(\mathbf{k}, t) a_\nu^*(\mathbf{k}, t) \\ &= \lim_{\Lambda \rightarrow \infty} \left[m_0 u^\mu(t) + q A^\mu(z, \Lambda) + \frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} k^\mu a^\nu(\mathbf{k}, t) a_\nu^*(\mathbf{k}, t) \theta(\Lambda + k_\lambda u^\lambda(t)) \right], \end{aligned}$$

with $m_0 = m - (q^2/4\pi^2) \Lambda$. We obtain therefore a result very similar to that discussed in Sec. III D: when a proper cut-off is chosen and only fields that already obey Maxwell equations are considered, the method of the $\mathcal{R}$ -integrals leads to expressions that formally correspond to the introduction of an infinite renormalization of the particle's mass. The analogy with the situation occurring with covariant regularization methods in quantum field theory is evident. It will be analyzed more closely in the following section.

VI. THE COMPARISON WITH QUANTUM ELECTRODYNAMICS

In the previous sections of this paper we have provided a complete diverge-free formulation of the classical electrodynamics of point charges. In the light of this result it is of course extremely interesting to reconsider the problem of the relationship between classical and quantum theories. It is in fact well known that the difficulties related to ultraviolet divergences in quantum electrodynamics have been up to now circumvented only in the framework of a perturbation expansion in the coupling constant $\alpha = e^2/4\pi\hbar c \simeq 1/137$. Finite non-perturbative expressions for the equations of motion or the hamiltonian operator have not yet been established, and it is even doubtful that they actually exist within the boundaries of pure electrodynamics. In connection with this problem, the implications of our new way of formulating the classical theory can be directly investigated by making use of the correspondence principle. Of course, in the quantum theory one encounters also the class of divergences associated with vacuum polarization, that have no direct equivalent in the classical theory. We shall not here take them into consideration, and we shall limit our analysis to the simplest situation in which the analogy with the classical case can be significant, i.e. the electron self-energy to the lowest order in α . We shall assume that the charged particles are represented by spin 1/2 fermions (electrons), and therefore described as the quanta of a Dirac field ψ . We shall follow the scheme of canonical quantization and, by analogy with the classical Poisson brackets (4.15) and (4.14), we shall postulate the usual fundamental equal-time commutation or anticommutation relations between field operators.

Considering the usual Gell-Mann and Low perturbative expansion of the two-point fermionic Green function $iS'_F(x) = \langle 0|T [\psi(x)\bar{\psi}(0)]|0\rangle$, the contribution of order α (using natural units with $c = \hbar = 1$) can be expressed as

$$iS_F^{(2)}(x) = \frac{(-i)^2}{2} \int_{-\infty}^{+\infty} dt_1 \int_{-\infty}^{+\infty} dt_2 \langle 0|T [H_I(t_1)H_I(t_2)\psi(x)\bar{\psi}(0)]|0\rangle_{(0)}, \quad (6.1)$$

where the subscript (0) indicates that the matrix element involves the non-interacting vacuum state and field operators, and only connected Feynman diagrams have to be taken into account. By analogy with Eq. (4.7) or (5.10), the interaction hamiltonian is obtained by substituting the field $\tilde{A}^\mu$ in place of A^μ into the familiar expression of the non-renormalized theory. This gives

$$H_I(t) = - \int d^3\mathbf{x} \tilde{A}_\mu(x) j^\mu(x) = -ie \int d^3\mathbf{x} \tilde{A}_\mu : \bar{\psi}\gamma^\mu\psi :, \quad (6.2)$$

where $\tilde{A}^\mu$ is given by a $\mathcal{R}$ -integral of the form (5.11). The natural way to generalize our notion of $\mathcal{R}$ -integral when the integrand is an operator in a Hilbert space is to perform the limiting procedure in the weak sense, i.e. on the matrix elements of the operator between any pair of states. This directly provides the prescription for calculating $iS_F^{(2)}$. One first has to replace the field $\tilde{A}_\mu(x)$ in Eq. (6.2) with the expression

$$A_\mu(x, \Lambda) = \frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} [a_\mu(\mathbf{k})e^{ik\cdot x} + a_\mu^*(\mathbf{k})e^{-ik\cdot x}] \theta(\Lambda - f(\mathbf{k})), \quad (6.3)$$

where $f(\mathbf{k})$ is an appropriate function of $\mathbf{k}$. Then, after taking the expectation value on the non-interacting vacuum state of the time-ordered product indicated in Eq. (6.1), one has to

evaluate a properly defined $\mathcal{R}$ -limit of the resulting expression for $\Lambda \rightarrow \infty$. One may notice the direct correspondence between Eq. (6.3) and Eq. (5.14) when putting $f(\mathbf{k}) = -k_\lambda u^\lambda$ and considering that, since the field operators in Eq. (6.1) satisfy the free-field equations, one can make the substitution

$$a_\mu(\mathbf{k}, t) \rightarrow a_\mu(\mathbf{k}) e^{-ik^0 t}.$$

It is easy to see that as a result of these prescriptions the expectation value in Eq. (6.1) has to be calculated by making use of the following free propagators:

$$\begin{aligned} \langle 0|T[A^\mu(x, \Lambda)A^\nu(0, \Lambda)]|0\rangle_{(0)} &= ig^{\mu\nu} \int \frac{d^4 k}{(2\pi)^4} e^{ik \cdot x} D_F(k, \Lambda) \equiv ig^{\mu\nu} D_F(x, \Lambda) \\ \langle 0|T[\psi(x)\bar{\psi}(0)]|0\rangle_{(0)} &= i \int \frac{d^4 p}{(2\pi)^4} e^{ip \cdot x} S_F(p) \equiv iS_F(x), \end{aligned}$$

with

$$D_F(k, \Lambda) \equiv -\frac{1}{k^2 + \lambda^2 - i\varepsilon} \theta(\Lambda - f(\mathbf{k}))$$

and

$$S_F(p) = \frac{ip \cdot \gamma - m}{p^2 + m^2 - i\varepsilon}.$$

As usual, a photon mass term λ has been included to remove infrared divergences, but it is expected that the limit $\lambda \rightarrow 0$ can ultimately be taken for all physically significant quantities. By applying Wick's theorem one finds for the Fourier transform of Eq. (6.1):

$$S_F^{(2)}(p) \equiv \int d^4 x S_F^{(2)}(x) e^{-ip \cdot x} = S_F(p) \Sigma^{(2)}(p) S_F(p),$$

with

$$\Sigma^{(2)}(p) = \mathcal{R} \lim_{\Lambda \rightarrow \infty} \Sigma^{(2)}(p, \Lambda) \quad (6.4)$$

and

$$\Sigma^{(2)}(p, \Lambda) = i \frac{\alpha}{4\pi^3} \int d^4 k D_F(k, \Lambda) \gamma_\mu S_F(p - k) \gamma^\mu.$$

In the last equation, the integration in dk^0 can be performed by closing the integration path in the complex plane and applying the residue theorem. This gives

$$\begin{aligned} \Sigma^{(2)}(p, \Lambda) &= \frac{\alpha}{2\pi^2} \int d^3 \mathbf{k} \theta(\Lambda - f(\mathbf{k})) \left\{ \left[\frac{1}{k^0} \frac{i(p - k) \cdot \gamma + 2m}{(p - k)^2 + m^2} \right]_{k^0 = \sqrt{\mathbf{k}^2 + \lambda^2}} \right. \\ &\quad \left. + \left[\frac{1}{k^0 - p^0} \frac{i(p - k) \cdot \gamma + 2m}{k^2 + \lambda^2} \right]_{k^0 = p^0 + \sqrt{(\mathbf{k} - \mathbf{p})^2 + m^2}} \right\}. \end{aligned}$$

It can be easily checked that, at variance with the expressions (5.12) and (5.14) for the classical self-energy of a point charge, in the integrand of the above equation the leading

terms of order $|\mathbf{k}|^{-2}$ cancel each other, whereas the non-vanishing sum of the terms of order $|\mathbf{k}|^{-3}$ gives rise to a contribution of order $\alpha \ln \Lambda$ to the electron self-energy. Considering the case $p^2 < 0$, we shall initially assume to choose the reference frame for which $\mathbf{p} = 0$, $p^0 = \sqrt{-p^2}$. Taking $f(\mathbf{k}) = |\mathbf{k}|/\beta$, where β is an arbitrary parameter, the integrals can thus be easily evaluated. Then, by imposing the Lorentz covariance, one can generalize the result for arbitrary p . This provides

$$\Sigma^{(2)}(p, \Lambda) = \frac{\alpha}{2\pi} \left[m \left(3 \ln \frac{2\beta\Lambda}{m} - \frac{1}{2} \right) + (ip \cdot \gamma + m) \left(\ln \frac{2\beta\Lambda}{m} - 2 \ln \frac{m}{\lambda} + \frac{3}{2} \right) \right] + \Sigma^{c(2)}(p), \quad (6.5)$$

where $\Sigma^{c(2)}(p)$ is expressible as a power series containing second and higher powers of $(ip \cdot \gamma + m)$ and, as it can be easily checked, coincides with the finite one-loop electron self-energy provided by the usual renormalization theory. The form of Eq. (6.5) obviously suggests the generalization of the notion of $\mathcal{R}$ -limit which may be applied in Eq. (6.4): given a function $F(\Lambda)$ having a behavior for of the type

$$F(\Lambda) = c \ln \Lambda + d + O(\Lambda^{-1}) \quad \text{for } \Lambda \rightarrow \infty,$$

we shall define

$$\mathcal{R} \lim_{\Lambda \rightarrow \infty} F(\Lambda) = d.$$

We obtain in this way

$$\Sigma^{(2)}(p) = A^{(2)} + B^{(2)} (ip \cdot \gamma + m) + \Sigma^{c(2)}(p), \quad (6.6)$$

with

$$A^{(2)} = \frac{\alpha}{2\pi} m \left(3 \ln \frac{2\beta}{m} - \frac{1}{2} \right) \\ B^{(2)} = \frac{\alpha}{2\pi} \left(\ln \frac{2\beta}{m} - 2 \ln \frac{m}{\lambda} + \frac{3}{2} \right).$$

One can see from Eq. (6.6) that $A^{(2)}$ represents a finite electromagnetic mass of the electron. If m is identified with the physical mass, the above result amounts to introducing in the hamiltonian a bare mass $m_0 = m - A^{(2)} + O(\alpha^2)$. Besides, $B^{(2)}$ can be considered as the second-order contribution to a finite (apart from infrared divergences) fermion wave-function renormalization constant $Z_2 = 1 - B^{(2)} + O(\alpha^2)$. One may observe that the arbitrary finite parameter β only appears in the unobservable quantities m_0 and Z_2 : therefore it simply allows for a finite renormalization of the theory, once the ultraviolet divergences have been removed by the $\mathcal{R}$ -limit procedure. One can then conclude that with the present method the familiar results of perturbative QED at order α are actually recovered without the need of introducing divergent parameters in the hamiltonian.

Of course, in order to really establish on a firm basis the correspondence between classical and quantum electrodynamics, one should be able to obtain such a result to any order in perturbation theory. However, an important difference between the two theories is given by

the fact that in the quantum case the full electron self-energy contains an infinite number of contributions of any order in α . When a manifestly covariant cut-off Λ is used (as in the Pauli-Villars regularization scheme) the contribution of order α^n contains powers of $\ln(\Lambda/m)$ up to $\ln^n(\Lambda/m)$. Therefore the simple logarithmic dependence which we have found for the lowest-order contribution in Eq. (6.5) cannot be assumed to represent the behavior of the exact electron self-energy as a function of Λ . We have just seen how the form of this behavior (which in turn is strictly related to the large momentum behavior of the renormalized self-energy) affects the correct definition of the $\mathcal{R}$ -limits that may appear in the equations of motion and the hamiltonian operator of QED. These considerations therefore seem to indicate that the problem of explicitly providing a finite and non-perturbative mathematical formulation of quantum electrodynamics is in any case ultimately related to the investigation of the short-distance behavior of the theory.

APPENDIX A: ASYMPTOTIC EXPANSION OF THE FIELD NEAR A POINT CHARGE

We want to construct a covariant asymptotic expansion for the retarded and advanced electromagnetic potentials near a point $z(\tau)$ of the world-line of a particle with proper time τ . Such an expansion will have the form

$$A_{\text{adv/ret}}^\mu(z+x) \sim (A_{\text{adv/ret}}^\mu)_{-1}(x) + (A_{\text{adv/ret}}^\mu)_0(x) + (A_{\text{adv/ret}}^\mu)_1(x) + \dots, \quad (\text{A1})$$

where each term $(A_{\text{adv/ret}}^\mu)_j(x)$ is a homogeneous function of x of degree j , and contains the time derivatives $d^k z/d\tau^k$ up to the order $k = j + 2$.

Clearly, the following two requirements must be satisfied:

1. When fixing $\bar{x} \equiv z + x$, since the left-hand side of Eq. (A1) becomes of course independent of τ , the series on the right-hand side must as a whole have the same property.
2. For $x^2 = x_\nu x^\nu = 0$ and $x^0 < 0$ (resp. $x^0 > 0$), the well-known expression of the Liénard-Wiechert advanced (resp. retarded) potentials must be obtained:

$$\begin{aligned} A_{\text{adv}}^\mu(z+x) &= \frac{q}{4\pi} \frac{u^\mu}{x_\nu u^\nu} \quad \text{for } x^2 = 0, \ x^0 < 0 \\ A_{\text{ret}}^\mu(z+x) &= -\frac{q}{4\pi} \frac{u^\mu}{x_\nu u^\nu} \quad \text{for } x^2 = 0, \ x^0 > 0. \end{aligned}$$

Conversely, it is easy to see that if we manage to obtain a series satisfying the requirements 1 and 2, then Eq. (A1) will hold for arbitrary x . Let us introduce the scalar

$$y = \sqrt{x^2 + (x \cdot u)^2} \quad (\text{A2})$$

and the four-vector

$$\bar{y}^\mu = x^\mu + x_\lambda u^\lambda u^\mu. \quad (\text{A3})$$

We note immediately that

$$\bar{y}_\mu u^\mu = 0, \quad (\text{A4})$$

$$\bar{y}_\mu \bar{y}^\mu = \bar{y}_\mu x^\mu = x_\mu x^\mu + (x_\nu u^\nu)^2 = y^2. \quad (\text{A5})$$

If we put

$$(A_{\text{adv}}^\mu)_{-1}(x) = (A_{\text{ret}}^\mu)_{-1}(x) = \frac{q}{4\pi} \frac{u^\mu}{y} \equiv (A^\mu)_{-1}(x), \quad (\text{A6})$$

we get a term of degree -1 which clearly satisfies the requirement 2. Furthermore, for fixed $\bar{x}$ we have

$$\frac{dx^\mu}{d\tau} = \frac{d}{d\tau} [\bar{x}^\mu - z^\mu(\tau)] = -u^\mu(\tau),$$

$$\frac{dy}{d\tau} = \frac{\partial y}{\partial x^\nu} \frac{dx^\nu}{d\tau} + \frac{\partial y}{\partial u^\nu} \frac{du^\nu}{d\tau} = \frac{1}{y} \left(-\bar{y}_\nu u^\nu + x_\lambda u^\lambda x_\nu \dot{u}^\nu \right) = \frac{x \cdot u}{y} x \cdot \dot{u},$$

where the last equality follows from Eq. (A4). Therefore:

$$\frac{d}{d\tau} (A^\mu)_{-1}(x) = \frac{q}{4\pi} \left[\frac{\dot{u}^\mu}{y} - (x \cdot u)(x \cdot \dot{u}) \frac{u^\mu}{y^3} \right]. \quad (\text{A7})$$

In order that the property 1 be satisfied, $(A_{\text{adv/ret}}^\mu)_0(x)$ must give rise, when differentiated with respect to τ , to a term of order x^{-1} that exactly cancels that given by Eq. (A7). Furthermore, because of requirement 2, we must have $(A_{\text{adv}}^\mu)_0(x) = 0$ for $x^2 = 0$, $x^0 < 0$, and $(A_{\text{ret}}^\mu)_0(x) = 0$ for $x^2 = 0$, $x^0 > 0$. It is straightforward to verify that these conditions are fulfilled by

$$(A_{\text{adv/ret}}^\mu)_0(x) = (A^\mu)_0(x) \pm \frac{q}{4\pi} \dot{u}^\mu, \quad (\text{A8})$$

with

$$\frac{4\pi}{q} (A^\mu)_0(x) = -\frac{u \cdot x}{y} \dot{u}^\mu - \frac{x^2}{2y^3} (\dot{u} \cdot x) u^\mu \quad (\text{A9})$$

(in formulae like (A8), the upper and lower signs of the symbols $\pm$ and $\mp$ refer to the advanced and retarded fields respectively). One can then iterate the procedure in order to obtain the higher order terms. In particular, the term of degree 1 is found to be

$$(A_{\text{adv/ret}}^\mu)_1(x) = (A^\mu)_1(x) \mp \frac{q}{4\pi} \left[(\ddot{u} \cdot x - \dot{u}^2 u \cdot x) \frac{u^\mu}{3} + (\dot{u} \cdot x) \dot{u}^\mu + (u \cdot x) \ddot{u}^\mu \right],$$

with

$$\begin{aligned} \frac{4\pi}{q} (A^\mu)_1(x) = & \left[\frac{3x^4}{8y^5} (\dot{u} \cdot x)^2 + \left(1 + \frac{x^2}{2y^2} \right) \frac{u \cdot x}{3y} \ddot{u} \cdot x - \left(2 - \frac{x^2}{y^2} - \frac{x^4}{4y^4} \right) \frac{y}{6} \dot{u}^2 \right] u^\mu \\ & + \left(1 + \frac{x^2}{2y^2} \right) \frac{u \cdot x}{y} (\dot{u} \cdot x) \dot{u}^\mu + \left(1 - \frac{x^2}{2y^2} \right) y \ddot{u}^\mu. \end{aligned}$$

From Eq. (A1) one can directly obtain, using Eq. (3.1), an analogous expansion for the advanced and retarded fields:

$$F_{\text{adv/ret}}^{\mu\nu}(z+x) = (F_{\text{adv/ret}}^{\mu\nu})_{-2}(x) + (F_{\text{adv/ret}}^{\mu\nu})_{-1}(x) + (F_{\text{adv/ret}}^{\mu\nu})_0(x) + \dots,$$

with

$$(F_{\text{adv/ret}}^{\mu\nu})_{j-1}(x) = \partial^\mu (A_{\text{adv/ret}}^\nu)_j(x) - \partial^\nu (A_{\text{adv/ret}}^\mu)_j(x).$$

This provides

$$(F_{\text{adv}})_{-2}(x) = (F_{\text{ret}})_{-2}(x) = \frac{q}{4\pi y^3} u \wedge x \equiv (F)_{-2}(x), \quad (\text{A10})$$

$$\begin{aligned} (F_{\text{adv}})_{-1}(x) &= (F_{\text{ret}})_{-1}(x) \\ &= \frac{q}{4\pi y^3} \left[\left(1 - \frac{3x^2}{2y^2}\right) (\dot{u} \cdot x) u \wedge x - (u \cdot x) \dot{u} \wedge x - \frac{x^2}{2} u \wedge \dot{u} \right] \equiv (F)_{-1}(x) \end{aligned} \quad (\text{A11})$$

and

$$(F_{\text{adv/ret}}^{\mu\nu})_0(x) = (F^{\mu\nu})_0(x) \mp \frac{2}{3} \frac{q}{4\pi} (u^\mu \ddot{u}^\nu - u^\nu \ddot{u}^\mu), \quad (\text{A12})$$

with

$$\begin{aligned} \frac{4\pi}{q} (F)_0 &= \frac{3x^2}{2y^5} (\dot{u} \cdot x) \left[\frac{x^2}{2} u \wedge \dot{u} + (u \cdot x) \dot{u} \wedge x \right] \\ &\quad + \frac{x^2}{2y^5} \left[3 \left(\frac{5x^2}{4y^2} - 1 \right) (\dot{u} \cdot x)^2 + \frac{x^2}{4} \dot{u}^2 + (u \cdot x) (\ddot{u} \cdot x) \right] u \wedge x \\ &\quad - \frac{x^2}{2y^3} \ddot{u} \wedge x + \frac{1}{3y} \left(2 + \frac{x^2}{y^2} \right) (u \cdot x) u \wedge \ddot{u}. \end{aligned} \quad (\text{A13})$$

APPENDIX B: DIVERGENT TERMS OF THE NON-RENORMALIZED ELECTROMAGNETIC ENERGY-MOMENTUM

Let us introduce, on the space at fixed t for the laboratory system, the spatial coordinates $\mathbf{y}_n$ corresponding to a reference system moving with the instantaneous velocity of the n -th particle:

$$\mathbf{y}_n = \mathbf{x} - \mathbf{z}_n + (\gamma_n - 1) \frac{(\mathbf{x} - \mathbf{z}_n) \cdot \mathbf{v}_n}{\mathbf{v}_n^2} \mathbf{v}_n. \quad (\text{B1})$$

The modulus of the vector $\mathbf{y}_n$ is equal to the scalar y_n given by Eq. (2.14), so that we can write $y_n = |\mathbf{y}_n|$. Inverting Eq. (B1), performing the translation $\mathbf{x} - \mathbf{z}_n \rightarrow \mathbf{x}$ and dropping the index n , one gets

$$\mathbf{x} = \mathbf{y} + \left(\frac{1}{\gamma} - 1 \right) \frac{\mathbf{y} \cdot \mathbf{v}}{\mathbf{v}^2} \mathbf{v}, \quad (\text{B2})$$

from which it follows

$$\mathbf{x} \cdot \mathbf{v} = \frac{1}{\gamma} \mathbf{y} \cdot \mathbf{v}, \quad (\text{B3})$$

$$\mathbf{x}^2 = y^2 - (\mathbf{y} \cdot \mathbf{v})^2. \quad (\text{B4})$$

Besides, from Eqs. (B2) and (A3) one obtains for $x^0 = 0$

$$\bar{y}^i = y^i + (\gamma - 1) \frac{y_k v^k}{v^2} v^i. \quad (\text{B5})$$

Defining the solid angle $d\Omega_{\mathbf{y}}$ such that $y^2 d\Omega_{\mathbf{y}} dy = d^3 \mathbf{y} = \gamma d^3 \mathbf{x}$, one obtains immediately

$$\int_{y=r} \frac{d\Omega_{\mathbf{y}}}{4\pi} y_i y_j = \frac{\delta_{ij}}{3} r^2. \quad (\text{B6})$$

Then, from Eqs. (B2), (B5) and (B6), it is straightforward to derive that

$$\int_{y=r} \frac{d\Omega_{\mathbf{y}}}{4\pi} x_i \bar{y}_j = \frac{\delta_{ij}}{3} r^2. \quad (\text{B7})$$

Let us now apply the above formulae to evaluate the $1/r$ term of the expression (2.12). From Eqs. (2.7) and (2.9) it follows that the leading term of T^{0k} can be expressed as

$$(T^{0k})_{-4} = (F^0_j)_{-2} (F^{kj})_{-2} = \frac{q^2}{4\pi y^6} \gamma x_j (u^k \bar{y}^j - u^j \bar{y}^k),$$

where we have used the fact that $u \wedge x = u \wedge \bar{y}$. By using Eq. (B7) it is easy to obtain that the corresponding leading term of $\bar{P}_{\text{em}}^\mu(t, r)$ is given by

$$\int d^3 \mathbf{x} (T^{0k})_{-4} \theta(y - r) = \gamma^{-1} \int_r^\infty y^2 dy \int d\Omega_{\mathbf{y}} (T^{0k})_{-4} = \frac{q^2}{4\pi r} \frac{2}{3} u^k, \quad (\text{B8})$$

in accordance with Eq. (2.15). We have also

$$(F^{\mu\nu})_{-2} (F_{\mu\nu})_{-2} = \frac{2q^2}{(4\pi)^2 y^6} u_\mu \bar{y}_\nu (u^\mu \bar{y}^\nu - u^\nu \bar{y}^\mu) = -\frac{2q^2}{(4\pi)^2 y^4}, \quad (\text{B9})$$

$$(F^{0i})_{-2} (F^0_i)_{-2} = \frac{q^2}{(4\pi)^2 y^6} \gamma^2 \mathbf{x}^2, \quad (\text{B10})$$

so that, using Eqs. (2.7), (B4) and (B6),

$$\begin{aligned} \int d^3 \mathbf{x} (T^{00})_{-4} \theta(y - r) &= \gamma^{-1} \int_r^\infty y^2 dy \int d\Omega_{\mathbf{y}} (T^{00})_{-4} \\ &= \frac{q^2}{4\pi} \int_r^\infty \frac{dy}{y^4} \int \frac{d\Omega_{\mathbf{y}}}{4\pi} \left(\gamma y^2 - \gamma y_i v^i y_j v^j - \frac{y^2}{2\gamma} \right) \\ &= \frac{q^2}{4\pi r} \left(\gamma - \frac{\gamma}{3} \mathbf{v}^2 - \frac{1}{2\gamma} \right) = \frac{q^2}{4\pi r} \left(\frac{2}{3} \gamma - \frac{1}{6\gamma} \right), \end{aligned} \quad (\text{B11})$$

to be compared with Eq. (2.16).

APPENDIX C: PROOF OF THEOREM 1

Let us consider a volume V containing a single charged particle. Omitting the particle's index we can write

$$\begin{aligned} \frac{d}{dt} \overline{P}_{V,\text{em}}^\mu(t, r) &= \frac{d}{dt} \int_V d^3\mathbf{x} T^{\mu 0} \theta(y - r) = \int_V d^3\mathbf{x} \partial_0 T^{\mu 0} \theta(y - r) + \int_V d^3\mathbf{x} T^{\mu 0} \partial_0 \theta(y - r) \\ &= - \int_V d^3\mathbf{x} \partial_i T^{\mu i} \theta(y - r) + \int_V d^3\mathbf{x} T^{\mu 0} \partial_0 \theta(y - r) \\ &= -\Phi_V^\mu(t) + \int_V d^3\mathbf{x} T^{\mu\nu} \partial_\nu \theta(y - r) = -\Phi_V^\mu(t) + \int_V d^3\mathbf{x} T^{\mu\nu} \partial_\nu y \delta(y - r), \quad (\text{C1}) \end{aligned}$$

where

$$\Phi_V^\mu(t) \equiv \int_{\partial V} d\sigma n_i T^{i\mu}(\mathbf{x}, t)$$

and we have used the equation $\partial_\nu T^{\nu\mu}(x) = 0$, which as a consequence of Maxwell equations holds for every x outside the particle's world-line. From Eq. (2.14) one obtains

$$\partial_i y \equiv \frac{\partial y}{\partial x^i} = \frac{1}{y} \left[x_i - z_i + (x_j - z_j) u^j u_i \right], \quad (\text{C2})$$

$$\partial_0 y \equiv \frac{\partial y}{\partial t} \Big|_{\mathbf{x}} = \frac{\partial y}{\partial z^j} \frac{dz^j}{dt} + \frac{\partial y}{\partial u^j} \frac{du^j}{dt} = -\frac{\partial y}{\partial x^j} u^j \gamma^{-1} + \frac{1}{y} (x_k - z_k) u^k (x_j - z_j) \dot{u}^j \gamma^{-1}. \quad (\text{C3})$$

We can now perform the translation $\mathbf{x} - \mathbf{z} \rightarrow \mathbf{x}$ and use the same notation of App. A. Then, applying the definition (A3) with $x^0 = 0$, we can rewrite Eqs. (C2) and (C3) as

$$\partial_i y = \frac{\overline{y}_i}{y}, \quad (\text{C4})$$

$$\partial_0 y = \frac{1}{\gamma y} \left(-\overline{y}_j u^j + x_j u^j x_h \dot{u}^h \right). \quad (\text{C5})$$

Therefore, making in the last term of Eq. (C1) the change of integration variables (B2) we obtain

$$\begin{aligned} \frac{d}{dt} \overline{P}_{V,\text{em}}^\mu(t, r) + \Phi_V^\mu(t) &= \gamma^{-1} \int_0^\infty y^2 dy \int d\Omega_{\mathbf{y}} T^{\mu\nu} \partial_\nu y \delta(y - r) \\ &= r \gamma^{-1} \int_{y=r} d\Omega_{\mathbf{y}} \left(T^{\mu i} \overline{y}_i - T^{\mu 0} \overline{y}_j u^j \gamma^{-1} + T^{\mu 0} x_j u^j x_h \dot{u}^h \gamma^{-1} \right). \quad (\text{C6}) \end{aligned}$$

In order to evaluate the behavior of the above expression for $r \rightarrow 0$, it is useful to expand the field near the particle as in Eq. (2.29):

$$F(x) = (F)_{-2}(x) + (F)_{-1}(x) + (F)_0(x) + \tilde{F} + O(x),$$

where $(F)_{-2}$, $(F)_{-1}$ and $(F)_0$ are given respectively by Eqs. (A10), (A11) and (A13), while according to Eq. (2.30) $\tilde{F} \equiv \tilde{F}(0) = \overline{F}(0)$. Using Eq. (2.7) one can perform a similar

expansion of the integrand on the right-hand side of Eq. (C6). Then, since $(F)_{-2}$ and $(F)_0$ are odd under space inversion, while $(F)_{-1}$ is even, it is easy to see that many terms of this expansion are spatially odd, and therefore give no contribution to the integral. Retaining only the terms with even parity, we obtain

$$\frac{d}{dt}\overline{P}_{V,\text{em}}^\mu(t, r) + \Phi_V^\mu(t) = (W^\mu)_{-1}(t)r^{-1} + (W^\mu)_0(t) + O(r), \quad (\text{C7})$$

with

$$(W^\mu)_{-1}(t)r^{-1} = r\gamma^{-1} \int_{y=r} d\Omega_{\mathbf{y}} \left[(T^{\mu i})_{-3} \overline{y}_i - (T^{\mu 0})_{-3} \overline{y}_j u^j \gamma^{-1} + (T^{\mu 0})_{-4} x_j u^j x_h \dot{u}^h \gamma^{-1} \right], \quad (\text{C8})$$

$$(W^\mu)_0(t) = r\gamma^{-1} \int_{y=r} d\Omega_{\mathbf{y}} \left\{ \tilde{F}^{\mu\nu} (F^i{}_\nu)_{-2} + (F^{\mu\nu})_{-2} \tilde{F}^i{}_\nu + \frac{g^{\mu i}}{2} \tilde{F}^{\lambda\nu} (F_{\lambda\nu})_{-2} \right. \\ \left. - \left[\tilde{F}^{\mu\nu} (F^0{}_\nu)_{-2} + (F^{\mu\nu})_{-2} \tilde{F}^0{}_\nu + \frac{g^{\mu 0}}{2} \tilde{F}^{\lambda\nu} (F_{\lambda\nu})_{-2} \right] \gamma^{-1} u^i \right\} \overline{y}_i. \quad (\text{C9})$$

In Eq. (C8), $(T^{\mu\nu})_{-4}$ and $(T^{\mu\nu})_{-3}$ are respectively the homogeneous terms of degree -4 and -3 of the energy-momentum tensor, which can be simply expressed as combinations of $(F)_{-2}$ and $(F)_{-1}$. However, by comparing Eq. (C7) with Eqs. (2.15)–(2.16) we obtain directly

$$(W^k)_{-1}r^{-1} = \frac{q^2}{4\pi r} \frac{2}{3} \dot{u}^k \gamma^{-1}, \quad (\text{C10})$$

$$(W^0)_{-1}r^{-1} = \frac{q^2}{4\pi r} \left[\frac{2}{3} \dot{u}^0 \gamma^{-1} - \frac{1}{6} \frac{d(\gamma^{-1})}{dt} \right], \quad (\text{C11})$$

$$(W^\mu)_0 = \frac{d}{dt} P_{V,\text{em}}^\mu + \Phi_V^\mu. \quad (\text{C12})$$

Of course, Eqs. (C10)–(C11) can also be verified starting from Eq. (C8) and carrying out a straightforward although somewhat lengthy calculation, which may only serve as a consistency check for the whole procedure. The integrations on the right-hand side of Eq. (C9) can instead be done immediately using Eqs. (2.9) and (B7). Taking into account Eq. (C12) this provides

$$\frac{d}{dt} P_{V,\text{em}}^\mu + \Phi_V^\mu = -q \tilde{F}^{\mu\nu} u_\nu \gamma^{-1}. \quad (\text{C13})$$

The equation of motion for the charged particle can then be obtained by postulating that, in accordance with Eqs. (2.17) and (2.19),

$$\frac{d}{dt} (P_{V,\text{em}}^\mu + m u^\mu) = -\Phi_V^\mu.$$

Using Eq. (C13) this yields

$$m \dot{u}^\mu = q \tilde{F}^{\mu\nu} u_\nu,$$

which corresponds to Eq. (2.20).

Vice versa, let us consider a volume V containing an arbitrary number of particles, all obeying the equation of motion (2.20). Then, by decomposing V as the union of a due number of subsets each containing a single particle, and then applying to each of them the equation (C13), it is immediate to prove that Eq. (2.17) is satisfied.

APPENDIX D: PROOF OF THEOREM 2

1. The one-particle case

Let us consider a tube T of arbitrary shape in four-dimensional space-time, containing the world-line of a single particle of charge q . Let us suppose also that a given point $P(\bar{\tau})$ of this world-line, corresponding to a value $\bar{\tau}$ of the proper time, is assigned the coordinates $z^\mu = z'^\mu = 0$ in two inertial reference frames O and O' . We shall define π and π' as the three-dimensional spaces characterized by the equations $x^0 = 0$ and $x'^0 = 0$ respectively. Clearly both spaces intersect the world-line at the point $P(\bar{\tau})$. Let us also define the vector

$$J^\mu(x) \equiv T^{\mu\nu}(x)b_\nu,$$

where b is an arbitrary constant four-vector. From the local conservation of the electromagnetic energy-momentum tensor $T^{\mu\nu}$, it follows immediately that $\partial_\mu J^\mu = 0$ everywhere in the tube T except on the particle's world-line. For r much smaller than the diameter of the tube T , we have via a by-parts integration

$$\begin{aligned} 0 &= \int_T d^4x \partial_\mu J^\mu \theta(x^0) \theta(-x'^0) \theta(y-r) \\ &= - \int_{T \cap \pi} d^3\mathbf{x} J^0 \theta(-x'^0) \theta(y-r) + \int_{T \cap \pi'} d^3\mathbf{x}' J'^0 \theta(x^0) \theta(y-r) \\ &\quad + \int_{\partial T} d\sigma J^\mu n_\mu \theta(x^0) \theta(-x'^0) - \int_T d^4x \frac{J^\mu \bar{y}_\mu}{y} \theta(x^0) \theta(-x'^0) \delta(y-r), \end{aligned} \quad (D1)$$

where n is the unitary four-vector normal to the element $d\sigma$ of the tube's surface ∂T , y and $\bar{y}^\mu$ are again defined according to Eqs. (A2)–(A3), and we have used the relation $\partial_\mu y = \bar{y}_\mu/y$ (note that here, at variance with the situation discussed in App. C, partial derivatives with respect to time are performed keeping constant the position and velocity of the particle). Similarly we have

$$\begin{aligned} 0 &= \int_T d^4x \partial_\mu J^\mu \theta(-x^0) \theta(x'^0) \theta(y-r) \\ &= \int_{T \cap \pi} d^3\mathbf{x} J^0 \theta(x'^0) \theta(y-r) - \int_{T \cap \pi'} d^3\mathbf{x}' J'^0 \theta(-x^0) \theta(y-r) \\ &\quad + \int_{\partial T} d\sigma J^\mu n_\mu \theta(-x^0) \theta(x'^0) - \int_T d^4x \frac{J^\mu \bar{y}_\mu}{y} \theta(-x^0) \theta(x'^0) \delta(y-r). \end{aligned} \quad (D2)$$

By subtracting Eq. (D2) to Eq. (D1) we obtain

$$\begin{aligned} &\int_{T \cap \pi} d^3\mathbf{x} J^0 \theta(y-r) - \int_{T \cap \pi'} d^3\mathbf{x}' J'^0 \theta(y-r) \\ &= \int_{\partial T} d\sigma J^\mu n_\mu [\theta(x^0) \theta(-x'^0) - \theta(-x^0) \theta(x'^0)] \\ &\quad - \int_T d^4x \frac{J^\mu \bar{y}_\mu}{y} [\theta(x^0) \theta(-x'^0) - \theta(-x^0) \theta(x'^0)] \delta(y-r). \end{aligned} \quad (D3)$$

According to the arguments discussed in Sec. II, for $r \rightarrow 0$ the two terms on the left-hand side of Eq. (D3) can be expressed in the form

$$\int_{T \cap \pi} d^3 \mathbf{x} J^0 \theta(y-r) = b_\nu (P^\nu)_{-1} r^{-1} + b_\nu \mathcal{R} \int_{T \cap \pi} d^3 \mathbf{x} T^{0\nu} + O(r), \quad (\text{D4})$$

$$\int_{T \cap \pi'} d^3 \mathbf{x}' J'^0 \theta(y-r) = b'_\nu (P'^\nu)_{-1} r^{-1} + b'_\nu \mathcal{R} \int_{T \cap \pi'} d^3 \mathbf{x}' T'^{0\nu} + O(r), \quad (\text{D5})$$

where, in accordance with Eqs. (2.15) and (2.16), we have defined

$$(P^k)_{-1} \equiv \frac{q^2}{4\pi} \frac{2}{3} u^k \quad (\text{D6})$$

$$(P^0)_{-1} \equiv \frac{q^2}{4\pi} \left(\frac{2}{3} u^0 - \frac{1}{6} \gamma^{-1} \right), \quad (\text{D7})$$

and similarly for the primed quantities.

Let us now assume that the reference frame O is related to O' by a boost transformation bringing the particle instantaneously at rest at the proper time $\bar{\tau}$. We then have $\mathbf{v} = 0$, $y = |\mathbf{x}|$, $\bar{y}^0 = 0$, $\bar{y}^k = x^k$, $x'^0 = \gamma' (x^0 + \mathbf{v}' \cdot \mathbf{x})$, so that the last term on the right-hand side of Eq. (D3) can be rewritten as

$$- \int d^4 x \frac{J^\mu \bar{y}_\mu}{y} \left[\theta(x^0) \theta(-x'^0) - \theta(-x^0) \theta(x'^0) \right] \delta(y-r) = r \int_{|\mathbf{x}|=r} d\Omega_{\mathbf{x}} \int_{-\mathbf{x} \cdot \mathbf{v}'}^0 dx^0 x_i T^{i\nu} b_\nu. \quad (\text{D8})$$

Again, it is here useful to expand the energy-momentum tensor as a series of homogeneous terms of increasing degree in x . By applying Eqs. (2.7) and (2.9) with $u^k = 0$, $u^0 = 1$, one obtains

$$(T^{i0})_{-4} = 0$$

$$(T^{ij})_{-4} = \frac{q^2}{(4\pi)^2 |\mathbf{x}|^4} \left(\frac{1}{2} \delta^{ij} - \frac{x^i x^j}{\mathbf{x}^2} \right),$$

so that the $1/r$ term of the expression (D8) is given by

$$r \int_{|\mathbf{x}|=r} d\Omega_{\mathbf{x}} \int_{-\mathbf{x} \cdot \mathbf{v}'}^0 dx^0 x_i (T^{i\nu})_{-4} b_\nu = -\frac{q^2}{8\pi r^3} \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} (\mathbf{x} \cdot \mathbf{v}') (\mathbf{x} \cdot \mathbf{b}) = -\frac{q^2}{24\pi r} (\mathbf{v}' \cdot \mathbf{b}). \quad (\text{D9})$$

This result could also have been deduced directly from Eq. (D3). In fact, using Eqs. (D4)–(D7) and the relations $b'_\nu u'^\nu = b_\nu u^\nu = b_0$, $b'^0 = \gamma' (b^0 + \mathbf{v}' \cdot \mathbf{b})$, the term of order $1/r$ on the left-hand side of Eq. (D3) can be written as

$$b_\nu (P^\nu)_{-1} r^{-1} - b'_\nu (P'^\nu)_{-1} r^{-1} = \frac{q^2}{4\pi r} \left[\frac{2}{3} (b_\nu u^\nu - b'_\nu u'^\nu) - \frac{1}{6} (b_0 - \gamma'^{-1} b'_0) \right] = -\frac{q^2}{24\pi r} (\mathbf{v}' \cdot \mathbf{b}),$$

in accordance with Eq. (D9). Thus Eq. (D9) is strictly related with the non-covariance of the divergent part $(P^\nu)_{-1} r^{-1}$ of the non-renormalized electromagnetic energy-momentum.

Now, considering the zeroth order term of Eq. (D8), we have

$$r \int_{|\mathbf{x}|=r} d\Omega_{\mathbf{x}} \int_{-\mathbf{x} \cdot \mathbf{v}'}^0 dx^0 x_i (T^{i\nu})_{-3} b_\nu = 0.$$

This result follows immediately from the antisymmetry of $(T^{i\nu})_{-3}$ under the change of variables $x^\mu \rightarrow -x^\mu$. Therefore from Eqs. (D3)–(D5) we obtain

$$b_\nu \mathcal{R} \int_{T \cap \pi} d^3 \mathbf{x} T^{0\nu} - b'_\nu \mathcal{R} \int_{T \cap \pi'} d^3 \mathbf{x}' T'^{0\nu} = \int_{\partial T} d\sigma J^\mu n_\mu [\theta(x^0)\theta(-x'^0) - \theta(-x^0)\theta(x'^0)] . \quad (\text{D10})$$

If a single charged particle is present in the whole space, one can consider the limit of Eq. (D10) when the spatial dimensions of the tube T tend to infinity. Assuming as usual that $T^{\mu\nu}$ vanishes sufficiently fast at infinity, the integral over ∂T on the right-hand side will vanish, so that Eq. (D10) becomes

$$b_\nu P_{\text{em}}^\nu - b'_\nu P_{\text{em}}'^\nu = b_\nu \mathcal{R} \int_\pi d^3 \mathbf{x} T^{0\nu} - b'_\nu \mathcal{R} \int_{\pi'} d^3 \mathbf{x}' T'^{0\nu} = 0. \quad (\text{D11})$$

Since b was arbitrarily chosen, the above equation implies that P_{em}^ν transforms as a four-vector. One may notice that the particle's equation of motion has not been used up to now in this section. This means that, in the one-particle case, the Lorentz covariance of the electromagnetic energy-momentum four-vector defined by Eq. (2.22) is a consequence of the Maxwell equations alone, and is verified at any point of the world-line of the particle for any arbitrary particle trajectory.

We would like to point out that Eq. (D10) holds also in the more general case when in place of O one considers another arbitrary system of reference O'' for which $\mathbf{v}''(\bar{\tau}) \neq 0$. Assuming again that $z''^\mu(\bar{\tau}) = 0$, and calling π'' the space with equation $x''^0 = 0$, this means that we can write

$$b''_\nu \mathcal{R} \int_{T \cap \pi''} d^3 \mathbf{x}'' T''^{0\nu} - b'_\nu \mathcal{R} \int_{T \cap \pi'} d^3 \mathbf{x}' T'^{0\nu} = \int_{\partial T} d\sigma J^\mu n_\mu [\theta(x''^0)\theta(-x'^0) - \theta(-x''^0)\theta(x'^0)] . \quad (\text{D12})$$

To prove this, one has simply to write down the equivalent of Eq. (D10) with O'' in place of O' :

$$b_\nu \mathcal{R} \int_{T \cap \pi} d^3 \mathbf{x} T^{0\nu} - b''_\nu \mathcal{R} \int_{T \cap \pi''} d^3 \mathbf{x}'' T''^{0\nu} = \int_{\partial T} d\sigma J^\mu n_\mu [\theta(x^0)\theta(-x''^0) - \theta(-x^0)\theta(x''^0)] .$$

Then subtracting this equation from Eq. (D10) directly yields Eq. (D12), as it is easy to check. We are going to make use of Eq. (D12) for the analysis of the many-particle case in the following subsection.

2. The many-particle case

Let us now assume that N particles are present, with $N > 1$. One can prove that also in this case the total energy-momentum of the system, defined according to Eq. (2.21), transforms as a four-vector when particles and fields satisfy the equations of motion (2.20) and (2.24)–(2.27). To show this, let us divide the whole space-time into N tubes $T_1, T_2, \dots, T_N$ each containing the world-line of a single particle (clearly some tubes will extend spatially to infinity). Given two arbitrary inertial frames O and O' , the spaces π and π' (again identified by the equations $x^0 = 0$ and $x'^0 = 0$ respectively) will intersect the world-line of the generic n -th particle at two distinct points corresponding to the values τ and τ' of the proper time.

Calling $\bar{x}^0 \equiv x^0(\tau')$, we introduce also the space $\bar{\pi}$ characterized by the equation $x^0 = \bar{x}^0$. Repeating the same arguments of App. C we can write

$$\begin{aligned} & m_n u_n^\mu(\tau) + \mathcal{R} \int_{T_n \cap \pi} d^3 \mathbf{x} T^{\mu 0} - m_n u_n^\mu(\tau') - \mathcal{R} \int_{T_n \cap \bar{\pi}} d^3 \mathbf{x} T^{\mu 0} \\ &= \int_{\partial T_n} d\sigma n_\nu T^{\mu \nu} \left[\theta(x^0) \theta(\bar{x}^0 - x^0) - \theta(-x^0) \theta(x^0 - \bar{x}^0) \right] \end{aligned} \quad (\text{D13})$$

(note that the expression in square brackets on the right-hand side is written in such a way that the above formula holds in both the cases $\bar{x}^0 > 0$ and $\bar{x}^0 < 0$). Besides, rewriting Eq. (D12) for the n -th particle at the proper time τ' , we have

$$\begin{aligned} & b_\nu \mathcal{R} \int_{T_n \cap \bar{\pi}} d^3 \mathbf{x} T^{0\nu} - b'_\nu \mathcal{R} \int_{T_n \cap \pi'} d^3 \mathbf{x}' T'^{0\nu} \\ &= \int_{\partial T_n} d\sigma J^\mu n_\mu \left[\theta(x^0 - \bar{x}^0) \theta(-x'^0) - \theta(\bar{x}^0 - x^0) \theta(x'^0) \right]. \end{aligned} \quad (\text{D14})$$

Multiplying Eq. (D13) by b_μ and summing the result to Eq. (D14) we arrive at

$$\begin{aligned} & b_\nu \left[m_n u_n^\nu(\tau) + \mathcal{R} \int_{T_n \cap \pi} d^3 \mathbf{x} T^{0\nu} \right] - b'_\nu \left[m_n u_n'^\nu(\tau') + \mathcal{R} \int_{T_n \cap \pi'} d^3 \mathbf{x}' T'^{0\nu} \right] \\ &= \int_{\partial T_n} d\sigma J^\mu n_\mu \left[\theta(x^0) \theta(-x'^0) - \theta(-x^0) \theta(x'^0) \right]. \end{aligned}$$

Finally, let us sum over n . The integrals over the various ∂T_n will cancel out (neglecting as usual the contributions at infinity), so that we are left with

$$b_\nu P^\nu - b'_\nu P'^\nu = 0,$$

where the total momenta P^ν and P'^ν are defined according to Eq. (2.21) in the two reference frames O and O' respectively. Again, due to the arbitrariness of b , the above equation implies the Lorentz covariance of the total energy-momentum four-vector.

APPENDIX E: VARIATION OF THE LAGRANGIAN

We shall consider here a single charged particle, since the generalization of the results to an arbitrary number of particles is absolutely trivial. We observe preliminarily that, by differentiating Eq. (3.16) with respect to δz , δu , $\delta \dot{u}$ etc., it is possible to expand the variation δA of the potential in a similar way:

$$\delta A(z + x) = (\delta A)_{-2}(x) + (\delta A)_{-1}(x) + (\delta A)_0(x) + \delta \tilde{A}(z) + O(x). \quad (\text{E1})$$

As a function of x , $(\delta A)_j(x)$ is a homogeneous term of degree j which under the transformation $x \rightarrow -x$ is odd for $j = -2$, $j = 0$, and even for $j = -1$. Moreover $\delta \tilde{A}(z)$ can be expressed as

$$\delta \tilde{A}^\mu(z) = \mathcal{R} \lim_{r \rightarrow 0} \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} \delta A^\mu(\mathbf{z} + \mathbf{x}, t). \quad (\text{E2})$$

We have in particular, using the variables $\bar{y}^\mu$ defined by Eq. (A3),

$$(\delta A^\mu)_{-2}(x) = \frac{q}{4\pi y^3} u^\mu \bar{y}_i \delta z^i. \quad (\text{E3})$$

In expressions such this, it is always assumed that $\delta z^i \ll y$, since in the following we shall first evaluate the derivatives of $\bar{L}(r)$ for fixed r , and only at a later stage we shall consider their behavior for $r \rightarrow 0$ in order to obtain the corresponding derivatives of the renormalized lagrangian L .

Let us first consider the electromagnetic part of the lagrangian. Since the fields satisfy Maxwell equations, we can equivalently replace in Eq. (3.8) the factor $\theta(|\mathbf{x} - \mathbf{z}| - r)$ with $\theta(y - r)$, with y defined according to Eq. (2.14). As it is easy to see using Eqs. (2.8)–(2.10), Eq. (3.10) will remain valid, with only a modification in the coefficient $(L)_{-1}$, which plays no role at all in our calculations. Using this new definition of $\bar{L}_{\text{em}}$ we have

$$\delta \bar{L}_{\text{em}}(r) = - \int d^3 \mathbf{x} F^{\mu\nu} \partial_\mu \delta A_\nu \theta(y - r) - \frac{1}{4} \int d^3 \mathbf{x} F^{\mu\nu} F_{\mu\nu} \delta y \delta(y - r). \quad (\text{E4})$$

Integrating by parts, the first term on the right-hand side can be rewritten as

$$\begin{aligned} - \int d^3 \mathbf{x} F^{\mu\nu} \partial_\mu \delta A_\nu \theta(y - r) &= - \frac{d}{dx^0} \int d^3 \mathbf{x} F^{0j} \delta A_j \theta(y - r) \\ &\quad + \int d^3 \mathbf{x} \partial_\mu F^{\mu\nu} \delta A_\nu \theta(y - r) \\ &\quad + \int d^3 \mathbf{x} F^{\mu\nu} \delta A_\nu \partial_\mu y \delta(y - r). \end{aligned} \quad (\text{E5})$$

Using the expansions (2.29) and (E1) we can write for $r \rightarrow 0$

$$\int d^3 \mathbf{x} F^{0j} \delta A_j \theta(y - r) = \frac{H}{r} + \mathcal{R} \int d^3 \mathbf{x} F^{0j} \delta A_j + O(r) \quad (\text{E6})$$

where, according to Eqs. (2.9), (E3) and (B7),

$$\frac{H}{r} = \int d^3 \mathbf{x} (F^{0j})_{-2} (\delta A_j)_{-2} \theta(y - r) = \frac{q^2}{12\pi r} u_i \delta z^i.$$

The second term on the right-hand side of Eq. (E5) vanishes because of Eq. (3.14), while the third can be evaluated making use of Eqs. (C4) and (C5). After a shift $\mathbf{x} - \mathbf{z} \rightarrow \mathbf{x}$ in the integration variables, followed by the transformation (B2), we obtain

$$\begin{aligned} \int d^3 \mathbf{x} F^{\mu\nu} \delta A_\nu \partial_\mu y \delta(y - r) &= \frac{r}{\gamma} \int_{y=r} d\Omega_{\mathbf{y}} \left[F^{i\nu}(\mathbf{z} + \mathbf{x}) \delta A_\nu(\mathbf{z} + \mathbf{x}) \bar{y}_i \right. \\ &\quad \left. + F^{0i}(\mathbf{z} + \mathbf{x}) \delta A_i(\mathbf{z} + \mathbf{x}) \left(-\bar{y}_j v^j + x_j v^j x_k \dot{u}^k \right) \right]. \end{aligned} \quad (\text{E7})$$

We can now substitute into Eq. (E7) the expansions (2.29) and (E1), and then discard all those terms that have vanishing integrals due to their antisymmetry under space inversion. In this way we obtain

$$\begin{aligned} \int d^3 \mathbf{x} F^{\mu\nu} \delta A_\nu \partial_\mu y \delta(y - r) &= \frac{C}{r} + \frac{r}{\gamma} \int_{y=r} d\Omega_{\mathbf{y}} \left\{ \left[(F^{i\nu})_{-2} \delta \tilde{A}_\nu + \tilde{F}^{i\nu} (\delta A_\nu)_{-2} \right] \bar{y}_i \right. \\ &\quad \left. - \left[(F^{0i})_{-2} \delta \tilde{A}_i + \tilde{F}^{0i} (\delta A_i)_{-2} \right] \bar{y}_j v^j \right\} + O(r), \end{aligned}$$

where C is a term linearly dependent on δz and δu , which we do not need to evaluate explicitly. The integral on the right-hand side can be easily calculated by making use of Eqs. (2.9), (E3) and (B7). This gives

$$\begin{aligned}
& \int d^3\mathbf{x} F^{\mu\nu} \delta A_\nu \partial_\mu y \delta(y-r) \\
&= \frac{C}{r} + \frac{q}{\gamma r^2} \int_{y=r} \frac{d\Omega_{\mathbf{y}}}{4\pi} \left\{ \left[(u^i x^\nu - u^\nu x^i) \delta \tilde{A}_\nu + \tilde{F}^{i\nu} u_\nu (x_j + x_k u^k u_j) \delta z^j \right] \bar{y}_i \right. \\
&\quad \left. - \left[x^i \delta \tilde{A}_i + \tilde{F}^{0i} u_i (x_h + x_k u^k u_h) \delta z^h \gamma^{-1} \right] \bar{y}_j u^j \right\} + O(r) \\
&= \frac{C}{r} - \frac{q}{\gamma} \left(u^\nu \delta \tilde{A}_\nu + \frac{1}{3} \tilde{F}^{\mu i} u_\mu \delta z_i \right) + O(r). \tag{E8}
\end{aligned}$$

The second term on the right-hand side of Eq. (E4) can be evaluated with a very similar procedure, using the relation

$$\delta y = -\frac{1}{y} \left[x_i - z_i + (x_j - z_j) u^j u_i \right] \delta z^i + \frac{1}{y} (x_j - z_j) u^j (x_i - z_i) \delta u^i,$$

which follows from Eq. (2.13). We obtain

$$\begin{aligned}
-\frac{1}{4} \int d^3\mathbf{x} F^{\mu\nu} F_{\mu\nu} \delta y \delta(y-r) &= \frac{r}{4\gamma} \int_{y=r} d\Omega_{\mathbf{y}} F^{\mu\nu}(\mathbf{z} + \mathbf{x}) F_{\mu\nu}(\mathbf{z} + \mathbf{x}) (\bar{y}_i \delta z^i - x_j u^j x_i \delta u^i) \\
&= \frac{D}{r} + \frac{r}{2\gamma} \int_{y=r} d\Omega_{\mathbf{y}} (F^{\mu\nu})_{-2} \tilde{F}_{\mu\nu} \bar{y}_i \delta z^i + O(r) \\
&= \frac{D}{r} + \frac{q}{\gamma r^2} \int_{y=r} \frac{d\Omega_{\mathbf{y}}}{4\pi} u^\mu x^\nu \tilde{F}_{\mu\nu} \bar{y}_i \delta z^i + O(r) \\
&= \frac{D}{r} + \frac{q}{3\gamma} \tilde{F}^{\mu i} u_\mu \delta z_i + O(r). \tag{E9}
\end{aligned}$$

Therefore, combining Eqs. (E6), (E8) and (E9), we obtain

$$\delta L_{\text{em}} = \mathcal{R} \lim_{r \rightarrow 0} \delta \bar{L}_{\text{em}}(r) = -\frac{d}{dt} \left(\mathcal{R} \int d^3\mathbf{x} F^{0j} \delta A_j \right) - q\gamma^{-1} u^\nu \delta \tilde{A}_\nu. \tag{E10}$$

Considering now the interaction lagrangian, we have from Eq. (3.9)

$$\begin{aligned}
\delta \bar{L}_{\text{int}}(r) &= q\delta v^i \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} A_i(\mathbf{z} + \mathbf{x}, t) + q\gamma^{-1} u^\nu \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} \delta A_\nu(\mathbf{z} + \mathbf{x}, t) \\
&\quad + q\gamma^{-1} u^\nu \int_{|\mathbf{x}|=r} \frac{d\Omega_{\mathbf{x}}}{4\pi} \partial_i A_\nu(\mathbf{z} + \mathbf{x}, t) \delta z^i. \tag{E11}
\end{aligned}$$

Keeping in mind Eqs. (3.6), (3.20) and (E2) we can then write

$$\delta L_{\text{int}} = \mathcal{R} \lim_{r \rightarrow 0} \delta \bar{L}_{\text{int}}(r) = q\delta v^i \tilde{A}_i(z) + q\gamma^{-1} u^\nu \delta \tilde{A}_\nu(z) + q\gamma^{-1} u^\nu \partial_i \tilde{A}_\nu(z) \delta z^i. \tag{E12}$$

We have finally

$$\delta L_{\text{part}} = m u_i \delta v^i. \tag{E13}$$

Therefore, summing Eqs. (E10), (E12) and (E13), we obtain Eq. (3.22).

APPENDIX F: TRANSLATIONAL INVARIANCE AND CONSERVATION LAWS

Let us consider the Eq. (3.22), which has been derived making use only of the field equations. When also the equation of motion (3.24) for the particles is satisfied, the second expression in square brackets vanishes and we have (considering a single particle to simplify the notation)

$$\delta L = \frac{d}{dt} \left[\left(mu_i + q\tilde{A}_i(z) \right) \delta z^i - \mathcal{R} \int d^3\mathbf{x} F^{0j} \delta A_j \right]. \quad (\text{F1})$$

Let us now examine the space translation of the whole system by an infinitesimal length $d\varepsilon$ along the direction x^k :

$$\begin{aligned} \delta z^i &= \delta_k^i d\varepsilon, \\ \delta A^\mu(x) &= -\partial_k A^\mu(x) d\varepsilon. \end{aligned}$$

Substituting these expressions in Eq. (F1), the manifest invariance of the lagrangian allows us to write

$$0 = \frac{dL}{d\varepsilon} = \frac{d}{dt} \left[mu_k + q\tilde{A}_k(z) + \mathcal{R} \int d^3\mathbf{x} F^{0j} \partial_k A_j \right] = \frac{dP_k}{dt},$$

with

$$P_k = mu_k + q\tilde{A}_k(z) + \mathcal{R} \int d^3\mathbf{x} F^{0j} \partial_k A_j. \quad (\text{F2})$$

We are now going to show that the conserved quantity P_k is equal to the k -th component of the total momentum defined by Eqs. (2.21) and (2.22). We have in fact from Eq. (2.7)

$$\mathcal{R} \int d^3\mathbf{x} F^{0j} \partial_k A_j = \mathcal{R} \int d^3\mathbf{x} T^{0k} + \mathcal{R} \int d^3\mathbf{x} F^{0j} \partial_j A^k. \quad (\text{F3})$$

But

$$\begin{aligned} \int d^3\mathbf{x} F^{0j} \partial_j A^\mu \theta(y-r) &= - \int d^3\mathbf{x} F^{0j} A^\mu \partial_j \theta(y-r) = - \int d^3\mathbf{x} F^{0j} A^\mu \frac{\bar{y}_j}{y} \theta(y-r) \\ &= -\frac{r}{\gamma} \int_{y=r} d\Omega_{\mathbf{y}} \bar{y}_j (F^{0j})_{-2} \left[(A^\mu)_{-1} + \tilde{A}^\mu(z) \right] + O(r) \\ &= -\frac{q}{r^2} \int_{y=r} \frac{d\Omega_{\mathbf{y}}}{4\pi} \bar{y}_j x^j \left[q \frac{u^\mu}{4\pi r} + \tilde{A}^\mu(z) \right] + O(r) \\ &= q^2 \frac{u^\mu}{4\pi r} - q\tilde{A}^\mu(z) + O(r), \end{aligned}$$

so that

$$\mathcal{R} \int d^3\mathbf{x} F^{0j} \partial_j A^\mu = -q\tilde{A}^\mu(z). \quad (\text{F4})$$

Therefore Eq. (F2) becomes

$$P^k = mu^k + \mathcal{R} \int d^3\mathbf{x} T^{0k},$$

in accordance with Eqs. (2.21) and (2.22).

Let us now instead consider an infinitesimal temporal translation of the system's evolution:

$$\begin{aligned}\delta z^i &= v^i dt, \\ \delta A^\mu(x) &= \partial_0 A^\mu(x) dt.\end{aligned}$$

Since the lagrangian L contains no explicit dependence on t , we obtain from Eq. (F1)

$$\frac{dL}{dt} = \frac{d}{dt} \left[\left(mu_i + q\tilde{A}_i(z) \right) v^i - \mathcal{R} \int d^3\mathbf{x} F^{0j} \partial_0 A_j \right],$$

or

$$\frac{dP^0}{dt} = 0,$$

with

$$\begin{aligned}P^0 &= \left(mu_i + q\tilde{A}_i(z) \right) v^i - \mathcal{R} \int d^3\mathbf{x} F^{0j} \partial_0 A_j - L \\ &= mu^0 + \mathcal{R} \int d^3\mathbf{x} T^{00} + q\tilde{A}^0(z) + \mathcal{R} \int d^3\mathbf{x} F^{0j} \partial_j A^0 = mu^0 + \mathcal{R} \int d^3\mathbf{x} T^{00},\end{aligned}\quad (\text{F5})$$

where the last equality follows again from Eq. (F4). Therefore P^0 is equal to the total energy that was introduced in Sec. II. In the form given by Eqs. (F2) and (F5), the four-vector P^μ can also be immediately identified with the canonical energy-momentum defined in Sec. IV. The derivation of the conservation laws carried out in the present section clearly represents a generalization of Nöther's theorem applied to the lagrangian defined by Eqs. (3.2)–(3.5).

APPENDIX G: LARGE WAVE-VECTOR EXPANSION OF THE FIELD GENERATED BY A POINT CHARGE

By inverting Eqs. (5.6)–(5.7) one obtains

$$(a^\mu)_j(\mathbf{k}, t) = \int d^3\mathbf{x} e^{-i\mathbf{k}\cdot\mathbf{x}} \left[k^0 (A^\mu)_j(x) + i (\partial_0 A^\mu)_{j-1}(x) \right], \quad (\text{G1})$$

where the particle index n has been omitted to simplify the notation. Therefore Eqs. (5.4)–(5.5) can be obtained in a straightforward way by substituting into Eq. (G1) the explicit expressions derived from Eqs. (A6) and (A9), and then carrying out the Fourier transformation. However, one can obtain Eqs. (5.3)–(5.5) in a shorter way by considering, for a given time $\bar{t}$, the field (analogous to the “Wentzel's field” of Ref. [5])

$$A_t^\mu(x) \equiv - \int d^4y D(x-y) j^\mu(y) \theta(\bar{t} - y^0), \quad (\text{G2})$$

where

$$j^\mu(y) = q \frac{u^\mu}{\gamma} \delta(\mathbf{y} - \mathbf{z}) = q \int_{-\infty}^{+\infty} d\tau u^\mu(\tau) \delta^4(y - z(\tau)) \quad (\text{G3})$$

is the current density carried by the particle, while D is the usual invariant function:

$$D(x) = -\frac{i}{(2\pi)^3} \int d^4k e^{ik \cdot x} \delta(k^2) \varepsilon(k^0) = \frac{1}{2\pi} \delta(x^2) \varepsilon(x^0). \quad (\text{G4})$$

It is easy to see that $A_t^\mu(x) = A_{\text{ret}}^\mu(x)$ for $[x - z(\bar{t})]^2 > 0$, so that we have in particular

$$A_t^\mu(\mathbf{x}, \bar{t}) = A_{\text{ret}}^\mu(\mathbf{x}, \bar{t}) \quad (\text{G5a})$$

$$\partial_0 A_t^\mu(\mathbf{x}, \bar{t}) = \partial_0 A_{\text{ret}}^\mu(\mathbf{x}, \bar{t}) \quad (\text{G5b})$$

for every $\mathbf{x}$. From Eqs. (G2)–(G4) we get

$$\begin{aligned} A_t^\mu(x) &= -q \int d^4y D(x - y) \int_{-\infty}^{\bar{\tau}} d\tau u^\mu(\tau) \delta^4(y - z(\tau)) \\ &= \frac{iq}{(2\pi)^3} \int d^4k \delta(k^2) \varepsilon(k^0) \int_{-\infty}^{\bar{\tau}} d\tau u^\mu(\tau) e^{ik \cdot [x - z(\tau)]}, \end{aligned} \quad (\text{G6})$$

where $\bar{\tau}$ is the particle's proper time for which $z^0(\bar{\tau}) = \bar{t}$. We can write for τ near $\bar{\tau}$

$$\begin{aligned} z(\tau) &= z + us + \frac{1}{2} \dot{u} s^2 + O(s^3) \\ u(\tau) &= u + \dot{u} s + O(s^2), \end{aligned}$$

where on the right-hand sides z , u and $\dot{u}$ stand for $z(\bar{\tau})$, $u(\bar{\tau})$ and $\dot{u}(\bar{\tau})$ respectively, while $s = \tau - \bar{\tau}$. Substituting the two above expressions into Eq. (G6) we obtain

$$\begin{aligned} A_t^\mu(x) &= \frac{iq}{(2\pi)^3} \int d^4k \delta(k^2) \varepsilon(k^0) e^{ik \cdot (x - z)} \\ &\quad \times \int_{-\infty}^0 ds e^{-ik \cdot us} \left[1 - ik \cdot \dot{u} \frac{s^2}{2} + O(s^3) \right] [u^\mu + \dot{u}^\mu s + O(s^2)] \\ &= \frac{iq}{(2\pi)^3} \int d^4k \delta(k^2) \varepsilon(k^0) e^{ik \cdot (x - z)} \left[i \frac{u^\mu}{k \cdot u} + \frac{\dot{u}^\mu}{(k \cdot u)^2} - u^\mu \frac{k \cdot \dot{u}}{(k \cdot u)^3} + O(|\mathbf{k}|^{-3}) \right] \\ &= -\frac{q}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} e^{ik \cdot (x - z)} \left[\frac{u^\mu}{k \cdot u} + i \frac{(k \cdot \dot{u})u^\mu - (k \cdot u)\dot{u}^\mu}{(k \cdot u)^3} + O(|\mathbf{k}|^{-3}) \right] + \text{c.c.} \end{aligned}$$

Keeping into account Eqs. (G5), a comparison of the above equation with Eqs. (5.1) provides directly Eqs. (5.3)–(5.5). Of course, it is implicitly assumed that the incident field $A_{\text{in}}^\mu \equiv A^\mu - \sum_n A_{n,\text{ret}}^\mu$ is sufficiently well-behaved, so that its contribution to $a^\mu(\mathbf{k}, t)$ for large $\mathbf{k}$ is at most of order $|\mathbf{k}|^{-3}$.

APPENDIX H: PROOF OF THEOREM 4

Taking into account Eqs. (4.7) and (4.16), it is clear that Eqs. (5.9)–(5.10) are equivalent to the equation

$$\mathcal{R} \int d^3\mathbf{x} \Gamma_{\text{em}}^\mu(A, \pi) = \frac{1}{2(2\pi)^3} \mathcal{R} \int \frac{d^3\mathbf{k}}{k^0} k^\mu a^\nu(\mathbf{k}, t) a_\nu^*(\mathbf{k}, t), \quad (\text{H1})$$

with

$$\begin{aligned}\Gamma_{\text{em}}^0(A, \pi) &\equiv \frac{1}{4} F_{ij} F^{ij} + \frac{1}{2} \pi_\mu \pi^\mu - \pi^i \partial_i A^0 + \pi^0 \partial_i A^i \\ &= \frac{1}{2} \left[\partial_0 A_\mu \partial_0 A^\mu + \partial_i A_\mu \partial_i A^\mu + \partial_i \left(A^i \partial_j A^j - A^j \partial_j A^i \right) \right],\end{aligned}\quad (\text{H2})$$

$$\Gamma_{\text{em}}^k(A, \pi) \equiv -\pi_\mu \partial^k A^\mu = \partial^0 A_\mu \partial^k A^\mu - \partial_j \left(A^0 \partial^k A^j \right) + \partial^k \left(A^0 \partial_j A^j \right). \quad (\text{H3})$$

Eq. (H1) can be rewritten as

$$\mathcal{R} \lim_{r \rightarrow 0} \Pi_{\text{em}}^\mu(t, r) = \mathcal{R} \lim_{\Lambda \rightarrow \infty} K_{\text{em}}^\mu(t, \Lambda), \quad (\text{H4})$$

with

$$\Pi_{\text{em}}^\mu(t, r) \equiv \int d^3 \mathbf{x} \Gamma_{\text{em}}^\mu(A, \pi) \prod_{n=1}^N \theta(y_n - r) \quad (\text{H5})$$

and

$$K_{\text{em}}^\mu(t, \Lambda) \equiv \frac{1}{2(2\pi)^3} \int \frac{d^3 \mathbf{k}}{k^0} k^\mu a^\nu(\mathbf{k}, t) a_\nu^*(\mathbf{k}, t) \theta(\Lambda - k^0). \quad (\text{H6})$$

In order to evaluate the two $\mathcal{R}$ -limits in Eq. (H4), it is convenient to decompose the field in the following way:

$$\partial_\nu A^\mu = \sum_{n=1}^N (\partial_\nu A_n^\mu)_{-2} + (\partial_\nu A^\mu)_{\text{res}} \quad (\text{H7})$$

$$a^\mu = \sum_{n=1}^N (a_n^\mu)_{-1} e^{-i\mathbf{k} \cdot \mathbf{z}_n} + (a^\mu)_{\text{res}}. \quad (\text{H8})$$

The two above equations provide the definition for $(\partial_\nu A^\mu)_{\text{res}}$ and $(a^\mu)_{\text{res}}$, which are of course related to each other by an equation similar to (5.7):

$$(\partial_\nu A^\mu)_{\text{res}}(x) = \frac{i}{2(2\pi)^3} \int \frac{d^3 \mathbf{k}}{k^0} k_\nu \left[(a^\mu)_{\text{res}}(\mathbf{k}, t) e^{i\mathbf{k} \cdot \mathbf{x}} - (a^\mu)_{\text{res}}^*(\mathbf{k}, t) e^{-i\mathbf{k} \cdot \mathbf{x}} \right]. \quad (\text{H9})$$

Let us first consider the expression (H6). Taking into account Eq. (H8), we have for $\Lambda \rightarrow \infty$

$$K_{\text{em}}^\mu = C^\mu \Lambda + D^\mu + O(\Lambda^{-1}), \quad (\text{H10})$$

with

$$2(2\pi)^3 C^\mu \Lambda = \sum_{n=1}^N \int \frac{d^3 \mathbf{k}}{k^0} k^\mu (a_n^\nu)_{-1} (a_{n,\nu})_{-1}^* \theta(\Lambda - k^0)$$

and

$$\begin{aligned}
2(2\pi)^3 D^\mu &= \sum_{n \neq m} \int \frac{d^3 \mathbf{k}}{k^0} k^\mu (a_n^\nu)_{-1} (a_{m,\nu})_{-1}^* e^{-i\mathbf{k} \cdot (\mathbf{z}_n - \mathbf{z}_m)} \\
&+ \sum_{n=1}^N \int \frac{d^3 \mathbf{k}}{k^0} k^\mu \left[(a_n^\nu)_{-1} (a_\nu)_{\text{res}}^* e^{-i\mathbf{k} \cdot \mathbf{z}_n} + (a^\nu)_{\text{res}} (a_{n,\nu})_{-1}^* e^{i\mathbf{k} \cdot \mathbf{z}_n} \right] \\
&+ \int \frac{d^3 \mathbf{k}}{k^0} k^\mu (a^\nu)_{\text{res}} (a_\nu)_{\text{res}}^* .
\end{aligned} \tag{H11}$$

It is easy to check that the integrals appearing in the expression of D^μ are indeed convergent. One has to observe in this respect that, for $|\mathbf{k}| \rightarrow \infty$,

$$(a^\nu)_{\text{res}}(\mathbf{k}, t) = \sum_{n=1}^N (a_n^\nu)_0(\mathbf{k}, t) e^{-i\mathbf{k} \cdot \mathbf{z}_n} + O(|\mathbf{k}|^{-3})$$

and that, as a result of Eqs. (5.4)–(5.5),

$$(a_n^\nu)_{-1} (a_{n,\nu})_0^* + (a_n^\nu)_0 (a_{n,\nu})_{-1}^* = 0. \tag{H12}$$

Considering now Eq. (H5), we shall explicitly consider only the temporal component, since the other three can be treated in exactly the same way. We first observe that the three-divergence on the right-hand side of Eq. (H2) gives no contribution to $\Pi_{\text{em}}^0(t, r)$ for vanishing r . In fact, assuming as usual that the fields vanish sufficiently fast at infinity, we have

$$\begin{aligned}
&\frac{1}{2} \int d^3 \mathbf{x} \partial_i (A^i \partial_j A^j - A^j \partial_j A^i) \prod_{n=1}^N \theta(y_n - r) \\
&= -\frac{1}{2r} \sum_{n=1}^N \int d^3 \mathbf{x} (A^i \partial_j A^j - A^j \partial_j A^i) \bar{y}_{n,i} \delta(y_n - r).
\end{aligned}$$

Each of the N terms of the sum for $n = 1, \dots, N$ on the right-hand side of the above equation has the form

$$-\frac{r}{2\gamma} \int_{y=r} d\Omega_{\mathbf{y}} (A^i \partial_j A^j - A^j \partial_j A^i) \bar{y}_i ,$$

where the index n has been dropped and a transformation of the space coordinates of the form (B2) has been carried out. Using the asymptotic expansions (3.16) and (3.19), with $(A^\mu)_{-1} = qu^\mu/4\pi y$ and $(\partial_\nu A^\mu)_{-2} = q\bar{y}_\nu u^\mu/4\pi y^3$, in the limit $r \rightarrow 0$ the above integral becomes

$$\begin{aligned}
&-\frac{r}{2\gamma} \int_{y=r} d\Omega_{\mathbf{y}} \left\{ [(A^i)_{-1} + \tilde{A}^i] (\partial_j A^j)_{-2} - [(A^j)_{-1} + \tilde{A}^j] (\partial_j A^i)_{-2} \right\} \bar{y}_i \\
&= \frac{r}{2\gamma} \int_{y=r} d\Omega_{\mathbf{y}} \frac{q}{4\pi r^3} \left\{ \left[\frac{qu^i}{4\pi r} + \tilde{A}^i \right] \bar{y}_j u^j - \left[\frac{qu^j}{4\pi r} + \tilde{A}^j \right] \bar{y}_j u^i \right\} \bar{y}_i = 0.
\end{aligned}$$

Therefore, by making use of Eq. (H7), we obtain that the behavior of $\Pi_{\text{em}}^0(t, r)$ for $r \rightarrow 0$ has the form

$$\Pi_{\text{em}}^0 = \frac{C'^0}{r} + D'^0 + O(r), \tag{H13}$$

with

$$\begin{aligned}
\frac{C'^0}{r} &= \frac{1}{2} \sum_{n=1}^N \int d^3\mathbf{x} \left[(\partial_0 A_{n,\mu})_{-2} (\partial_0 A_n^\mu)_{-2} + (\partial_i A_{n,\mu})_{-2} (\partial_i A_n^\mu)_{-2} \right] \theta(y_n - r), \\
D'^0 &= \sum_{n < m} \int d^3\mathbf{x} \left[(\partial_0 A_{n,\mu})_{-2} (\partial_0 A_m^\mu)_{-2} + (\partial_i A_{n,\mu})_{-2} (\partial_i A_m^\mu)_{-2} \right] \\
&\quad + \sum_{n=1}^N \int d^3\mathbf{x} \left[(\partial_0 A_\mu)_{\text{res}} (\partial_0 A_n^\mu)_{-2} + (\partial_i A_\mu)_{\text{res}} (\partial_i A_n^\mu)_{-2} \right] \\
&\quad + \frac{1}{2} \int d^3\mathbf{x} \left[(\partial_0 A_\mu)_{\text{res}} (\partial_0 A^\mu)_{\text{res}} + (\partial_i A_\mu)_{\text{res}} (\partial_i A^\mu)_{\text{res}} \right]. \tag{H14}
\end{aligned}$$

The fact that D'^0 is a finite quantity can be as usual verified using the asymptotic expansion (3.19) and observing that

$$\int d^3\mathbf{x} (\partial_\nu A_{n,\mu})_{-2} (\partial_\lambda A_n^\mu)_{-1} \theta(y_n - r) = 0 \tag{H15}$$

because of the spatial antisymmetry of the integrand. It may be noted that Eqs. (H12) and (H15) are directly related to each other. They are respectively the reasons why there is not a term of order $\ln r$ on the right-hand side of Eq. (H13) and a term of order $\ln \Lambda$ on the right-hand side of Eq. (H10).

Finally, by substituting Eqs. (5.7) and (H9) into Eq. (H14), carrying out the integrations over the $\mathbf{x}$ space and comparing the result with the right-hand side of Eq. (H11), it is straightforward to verify that $D^0 = D'^0$. Keeping in mind the definition of the $\mathcal{R}$ -limits, the Eq. (H4) is therefore proved.

To complete the demonstration of the theorem, we are now going to prove Eq. (5.11). We first observe that, according to Eqs. (3.16) and (3.17), we have

$$\tilde{A}^\mu(z_n) = \lim_{x \rightarrow 0} \left[A^\mu(z_n + x) - (A_n^\mu)_{-1}(x) - (A_n^\mu)_0(x) \right],$$

so that, taking into account Eq. (5.6), we get

$$\tilde{A}^\mu(z_n) = \frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} \left[(a_n^\mu)_R(\mathbf{k}, t) e^{i\mathbf{k} \cdot \mathbf{z}_n} + (a_n^\mu)_R^*(\mathbf{k}, t) e^{-i\mathbf{k} \cdot \mathbf{z}_n} \right], \tag{H16}$$

with

$$(a_n^\mu)_R \equiv a^\mu - (a_n^\mu)_{-1} e^{-i\mathbf{k} \cdot \mathbf{z}_n} - (a_n^\mu)_0 e^{-i\mathbf{k} \cdot \mathbf{z}_n}.$$

On the other hand, we have for $\Lambda \rightarrow \infty$

$$\frac{1}{2(2\pi)^3} \int \frac{d^3\mathbf{k}}{k^0} \left[a_\mu(\mathbf{k}, t) e^{i\mathbf{k} \cdot \mathbf{z}_n} + a_\mu^*(\mathbf{k}, t) e^{-i\mathbf{k} \cdot \mathbf{z}_n} \right] \theta(\Lambda - k^0) = E_\mu \Lambda + F_\mu + O(\Lambda^{-1}), \tag{H17}$$

with

$$\begin{aligned}
E^\mu \Lambda &= \frac{1}{2(2\pi)^3} \int \frac{d^3 \mathbf{k}}{k^0} \left[(a_n^\mu)_{-1}(\mathbf{k}, t) + (a_n^\mu)^*_{-1}(\mathbf{k}, t) \right] \theta(\Lambda - k^0) \\
&= \frac{1}{(2\pi)^3} \int \frac{d^3 \mathbf{k}}{k^0} (a_n^\mu)_{-1}(\mathbf{k}, t) \theta(\Lambda - k^0),
\end{aligned}$$

$$F^\mu = \frac{1}{2(2\pi)^3} \int \frac{d^3 \mathbf{k}}{k^0} \left[(a_n^\mu)_R(\mathbf{k}, t) e^{i\mathbf{k} \cdot \mathbf{z}_n} + (a_n^\mu)^*_R(\mathbf{k}, t) e^{-i\mathbf{k} \cdot \mathbf{z}_n} \right]. \quad (\text{H18})$$

One has to observe that $(a_n^\mu)_0$ does not contribute to Eq. (H17) since

$$(a_n^\mu)_0(\mathbf{k}, t) + (a_n^\mu)^*_0(\mathbf{k}, t) = 0.$$

This corresponds to the fact that $(A_n^\mu)_0$ is an antisymmetric function of $\mathbf{x}$, and so does not contribute to the integral on the right-hand side of Eq. (3.17). Taking into account Eq. (H17), the comparison of Eq. (H18) with Eq. (H16) provides finally Eq. (5.11).

REFERENCES

- [1] P. A. M. Dirac, *Proc. R. Soc. London, Ser. A* **167** (1938), 148.
- [2] F. Rohrlich, “Classical Charged Particles,” Addison-Wesley, Reading, MA, 1965; F. Rohrlich, in “The Physicist’s Conception of Nature” (J. Mehra, Ed.), Dordrecht-Holland, 1973.
- [3] A. O. Berut, “Electrodynamics and Classical Theory of Fields and Particles,” 2nd ed., Dover, New York, 1980; S. Coleman, in “Electromagnetism: Paths to Research” (D. Teplitz, Ed.), Plenum Press, New York, 1982; P. Pearle, *ibid.*
- [4] A. D. Fokker, *Z. Phys.* **58** (1929), 386.
- [5] P. A. M. Dirac, *Ann. Inst. Henry Poincaré* **9** (1939), 13.
- [6] F. Rohrlich, *Phys. Rev. Lett.* **12** (1964), 375.
- [7] C. Teitelboim, *Phys. Rev. D* **1** (1970), 1572; **3** (1971), 297; E. G. P. Rowe, *ibid.* **18** (1978), 3639.
- [8] E. G. P. Rowe, *Phys. Rev. Lett.* **53** (1984), 1779.
- [9] C. Teitelboim, *Phys. Rev. D* **4** (1971), 345.
- [10] C. Teitelboim and C. Lopez, *Lett. Nuovo Cimento* **2** (1971), 225.
- [11] F. E. Low, *Ann. Phys. (N.Y.)* **266** (1998), 274.